

For research use only. Not for clinical diagnosis.

Catalog No. BAM-03-005-5-EX

Keratinocyte Growth Factor (KGF/ FGF7), human, active

BACKGROUND

Keratinocyte Growth Factor (KGF), also known as Fibroblast Growth Factor 7 (FGF-7), is a member of fibroblast growth factor (FGF) family. Although FGF-7 has heparin binding activity similar to FGF-1, its mitogenic activity is predominantly exhibited in keratinocytes. It is not effective to fibroblasts and endothelial cells.

The human FGF-7 lacking the signal sequence (1-31 aa) was expressed in E. coli and purified by the chromatographic procedures. This product is an intact enzyme without tag with 19 kDa size (Fig.).

Applications: 1. Mitogen for epithelial cells

Western blot control for anti-FGF-7 antibodies
 Acceleration of wound healing is implied.
 Acceleration of hair development is implied.

Size: 5x 50 μg

Form: 1.0 mg / ml in PBS (10mM Na-phosphate, 150mM NaCl) pH7.2, 50% glycerol,

filter-sterilized

Purity: >95% as determined by SDS-PAGE (CBB staining)

Measurement of the

activity:

The ED50 as determined by a cell proliferation assay using MTS assay kit (CellTiter 96,

Promega) with human keratinocyte JCRB141cells was <10 ng/ml.

Storage: -20°C (long period, -80°C)

Data Link GenelD: 2252

Gene Sequence: M60828.1
Amino Acid Sequence: P21781

References: 1) Rubin JS et al. (1989) "Purification and characterization of a newly identified growth factor specific for

epithelial cells." Proc Natl Acad Sci USA 86: 802-806 PMID: 2915979

2) Aaronson SA et al. (1991) "Keratinocyte growth factor. A fibroblast growth factor family member with unusual

target cell specificity." Ann NY Acad Sci 638:62-77 PMID: 1664700




Fig. SDS-PAGE of human FGF-7

For research use only. Not for clinical diagnosis.

Manufactured by BioAcademia,Inc.



COSMO BIO CO., LTD.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

URL: http://www.cosmobio.co.jp e-mail: <u>export@cosmobio.co.jp</u> [回内連絡先] Phone: +81-3-5632-9610