

## Anti-GST antibody, rabbit serum

60-021, 100 ul

**Glutathione S transferase (GST)** from *Schistosoma japonicum* is commonly used to create fusion proteins. GST-tag has the size of 220 amino acids (roughly 26kDa) and is fused to the N-terminus of a protein. GST fusion proteins can be produced in *Escherichia coli*, as recombinant proteins and are used to purify and detect proteins of interest. The GST part binds its substrate, glutathione. GST-fusions protein can be easily purified from cell extracts by affinity chromatography with glutathione resin.

### Applications:

1. Western blotting (dilution: 1/2,000~1/10,000)
2. Immunoprecipitation (assay dependent)
3. ELISA

Other applications have not been tested.

**Immunogen:** Recombinant full-size GST (aa 1-212)

**Form:** Antiserum added with 0.05% sodium azide

**Specificity:** Specific to GST and GST-tagged proteins

**Storage:** Shipped at 4°C or -20°C, and upon arrival, aliquot and store at -20°C.

**Data Link:** NCBI Protein Data [AAA57089](http://www.ncbi.nlm.nih.gov/nuccore/AAA57089)

### References:

1. Smith DB & Johnson KS (1988) "Single-step purification of polypeptides expressed in *Escherichia coli* as fusions of glutathione-S-transferase." *Gene* **67**:31-40 PMID: [3047011](https://pubmed.ncbi.nlm.nih.gov/3047011/)
2. Kaelin WG Jr *et al* (1991) "Identification of cellular proteins that can interact specifically with the T/E1A-binding region of the retinoblastoma gene product." *Cell* **64**:521-532 PMID: [1825028](https://pubmed.ncbi.nlm.nih.gov/1825028/)
3. *Molecular Cloning: A Laboratory Manual* (eds. Sambrook, J., Russell, D.W. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, USA, 2001) pp.15.36-15.39, pp.18.48-18.59.

Fig.1 Detection of GST-tagged protein with this antibody by Western blotting.

- : Lysate of 293T cells transfected with an empty vector
- +: Lysate of 293T cells transfected with the plasmid carrying the GST-tagged importin gene

Fig.2 Immunoprecipitation of GST-tagged protein with this antibody followed by Western blotting.

- : Lysate of 293T cells transfected with an empty vector
- +: Lysate of 293T cells transfected with the plasmid carrying the GST-tagged INbox gene

Fig.1




Fig.2

