

POLYCLONAL ANTIBODY

For research use only. Not for clinical diagnosis.

Catalog No. BAM-62-010-EX

Anti-Sua7p/TFIIB (yeast)

BACKGROUND

The fundamental transcription factor TFIIB has the characteristics of stabilizing the DNA binding of TATA box-binding protein (TBP) and binding directly to DNA by its conformational change. Also its N terminal region binds to the RNA channel of RNA polymerase undertaking a very important role in the determination of transcription initiation point and promoter clearance. Sua7p is the TFIIB of budding yeast and is composed of 346 amino acid residues (aa).

The product was prepared by immunizing rabbit with recombinant protein which was over-expressed in E. coli with a plasmid carrying the entire Sua7p protein (1-346aa) of budding yeast, and purified by chromatography.

Using the product as antiserum in Western blotting, the band of 41 kD pertaining to Sua7p was obtained from the extract of yeast cells (Fig. 1).

Primary antibodies Product type

Rabbit Host Serum Source **Form** Liquid

0.1% sodium azide added to the antiserum

Volume 250 ul

Concentration **Specificity**

Recombinant protein which was over-expressed in E. coli with a plasmid carrying the Antigen

entire Sua7p protein (1-346aa) of budding yeast

Isotype

Application notes WB, ELISA

Recommended use

It can be used in Western blotting or ELISA for detection and titration of budding yeast

Recommended dilutions

Optimal dilutions/concentrations should be determined by the end user.

Data Link: SGD SUA7/YPR086W

Staining Pattern

Cross reactivity

Storage

4°C

1) Pinto I et al "The yeast SUA7 gene encodes a homolog of human transcription factor TFIIB and is required for References normal start site selection in vivo" Cell 68: 977-988 (1992) PMID: 1547497

> 2) Kornberg RD "Mechanism and regulation of yeast RNA polymerase II transcription" Cold Spring Harb Symp Quant Biol 63: 229-232 (1998) PMID: 10384286






Fig.1 Detection of Sua7p by Western blotting using the anti-Sua7p antibody.

Lane1, Extract of budding yeast
The antiserum was diluted 5000 fold before use.

For research use only. Not for clinical diagnosis.

Manufactured by BioAcademia,Inc.



COSMO BIO CO., LTD.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN