



Anti-Sds22 (*S. pombe*) antibody, rabbit serum

BACKGROUND

Schizosaccharomyces pombe **Sds22** protein contains leucine-rich repeats and physically interacts with the catalytic subunits of two type 1 protein phosphatases (Dis2 and Sds21). **Sds22** is a regulatory subunit of these phosphatases and the **Sds22**-bound phosphatases carry a key phosphatase activity essential for the progression from metaphase to anaphase. **Sds22** is essential for cell viability and in its absence, cells were blocked in metaphase. **Sds22** protein is predicted to form a repeating helical rod that is capable of enhancing a PP1-dependent dephosphorylation activity.

Product type	Primary antibodies
Host	Rabbit
Source	serum
Form	Liquid Rabbit antiserum added with 0.05 % sodium azide
Volume	100 µl
Concentration	
Specificity	Sds22
Antigen	Recombinant C-terminal region (1.8kb) of <i>S. pombe</i> Sds22 (1)
Isotype	

Application notes Immunoblotting, Immunoprecipitation, Immunofluorescence microscopy

Recommended use

Recommended dilutions

Immunoblotting (dilution: 1/200~1/500)

Optimal dilutions/concentrations should be determined by the end user.

Data Link: Swiss-Prot [P22194](#)

Staining Pattern

Cross reactivity Specific to *S. pombe*

Storage -20°C

References

- 1) Ohkura H and Yanagida M "S.pombe gene sds22+ essential for a midmitotic transition encodes a leucine-rich repeat protein that positively modulates protein phosphatase-1." *Cell* **64**: 149-157 (1991) PMID: [1846086](#)
- 2) Stone EM *et al* "Mitotic regulation of protein phosphatases by the fission yeast sds22 protein." *Curr Biol* **3**: 13-26 (1993) PMID: [15335873](#)
- 3) Ishii K *et al* "Requirement for PP1 phosphatase and 20S cyclosome/APC for the onset of anaphase is lessened by the dosage increase of a novel gene sds23+." *EMBO J.* **15**: 6629-6640 (1996) PMID: [8978689](#)




Fig.1, Immunoblot with anti-Sds22 antiserum of yeast extracts from (1) wild type strain HM123, (2) *sds22::ura4+* deletion mutant carrying pHR140-2 (ref.2).

The 40kD protein band was identified by immunoblot analysis of wild-type strain using anti-Sds22 antisera (lane1). The 40 kD band is enhanced in the *sds22::ura4+* disruption mutant strain that is rescued by the multicopy *sds22+* plasmid pHR140-2 (lane2).




Fig.2, Sds22 coprecipitates with Dis2 and Sds21 (ref.2).

Yeast extracts of wild type (wt) strain HM123 (lane 1 and 2), *dis2::ura4+* deletion mutant ($\Delta d2$, lane 3), *sds21::ura4+* deletion mutant ($\Delta s21$, lane 4) were immunoprecipitated followed by immunoblotting with the indicated antiserum, to detect the Sds22 or Dis2/Sds21 proteins.

Lane 1 was immunoprecipitated with the appropriate preimmune serum, lane 2-4 with the anti-Sds22 serum.

(a) denotes anti-Sds22 immunoblot; (b) denotes anti-D2C immunoblot. Anti-D2C crossreacts with both Sds21 and Dis2.

Anti-Sds22 antiserum coprecipitates both Dis2 and Sds21 proteins in the wild type strain (lane 2b). Consistently, Sds21 alone is precipitated in the *dis2* deletion mutant (lane 3b), and Dis2 alone is precipitated in the *sds21* deletion mutant (lane 4b).




Fig.3, Sds22 subcellular localization

Indirect immunofluorescence microscopy was performed by staining methanol fixed cells with (first column) anti-Sds22 antiserum, or (second column) DAPI to visualize chromosomal DNA.

(a) wild type HM123; (b) HM123 carrying multicopy *sds22+* plasmid pHR140-2.

Anti-Sds22 antibody stains the cytoplasm as well as the non-chromosomal domain of the nucleus of a wild type strain, as shown in (a). Nuclear staining increases in strains carrying a multicopy *sds22+* plasmid (b).

For research use only. Not for clinical diagnosis.

Manufactured by BioAcademia, Inc.



COSMO BIO Co., LTD.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

URL: <http://www.cosmobio.co.jp>

e-mail: export@cosmobio.co.jp

[Outside Japan] Phone : +81-3-5632-9617

[国内連絡先] Phone : +81-3-5632-9610

FAX : +81-3-5632-9618

FAX : +81-3-5632-9619