

MONOCLONAL ANTIBODY

For research use only. Not for clinical diagnosis.

Catalog No. 65-069EX

Anti-Hepatitis C Virus (HCV) NS5b protein (NS5B-6)FITC-labeled IgG

BACKGROUND

Hepatitis C virus (HCV) is a small (55-65 nm in size), enveloped, positive sense single-stranded RNA virus in the family Flaviviridae and the principal cause of parenteral non-A, non-B hepatitis. The virus genome consists of a single open reading frame of approximately 9,400 bases which encodes a single polyprotein of about 3,010 amino acids (1, 2, 3). The polyprotein is processed by host cell and viral proteases into four structural proteins (core, envelope1 and 2, and p7) and six non-structural proteins (NS2, 3, 4a, 4b, 5a, and 5b) necessary for viral replication. NS5b forms the C-terminal protein of the HCV polyprotein and is thought to represent the viral polymerase involved in RNA replication

This product is an FITC-labeled IgG ([FITC]/IgG = 6.0) produced from the IgG fraction.

Product type Primary antibodies

Host Mouse

Source

Purified monoclonal antibody (IgG) 1.3mg/ml in PBS, 50% glycerol, filter-sterilized **Form**

Volume 50µg

Concentration

Specificity Specific to human HCV NS5b protein of genotype 1b. Not tested in other genotypes. Antigen

A region of NS5a protein (the nucleotide sequence is shown in ref.4) of HCV

genotype 1b expressed in E.coli

NS5B-6 Clone

Mouse IgG2b kappa Isotype

Application notes

1.WB

2. Immunofluorescence staining 3. ELISA 4. FACS

Other applications have not been tested.

Optimal dilutions/concentrations should be determined by the end user.

UniProtKB HCV protein

Storage

References

(This antibody is produced and used ref.4)

Shipped at 4°C or with blue ice and store at -20°C

- 1. Brass V et al (2006) Molecular Virology of Hepatitis C Virus (HCV): 2006 Update. Int J Med Sci 3: 29-34 PMID: 16614739
- 2. Kato N et al (1990) "Molecular cloning of the human hepatitis C virus genome from Japanese patients with non-A, non-B hepatitis" Proc Natl Acad Sci USA 87: 9524-9528 PMID: 2175903
- 3. Takamizawa A et al (1991) "Structure and organization of the hepatitis C virus genome isolated from human carriers" J Virol 65: 1105-1113 PMID: 1847440
- 4. Manabe S et al (1994) "Production of nonstructural proteins of hepatitis C virus requires a putative viral protease encoded by N3" Virology 198: 636-644 PMID: 8291245

Related products

#65-066EX anti-HCV NS5b antibody

#65-068EX anti-HCV NS5b antibody biotinylated



Anti-Hepatitis C Virus (HCV) NS5b protein (NS5B-6)FITC-labeled




Fig.1 Western blotting of HCV NS5b protein.

Chimp liver cells were infected with recombinant vaccinia virus containing a HCV genome cDNA and were subjected to Western blotting using this antibody. The antibody detected as a 58-kDa band.



Fig. 2 Detection of HCV NS5b protein by immunofluorescence antibody staining.

Chimp liver cells were infected with recombinant
vaccinia virus containing a HCV genome cDNA.

After incubation for 48 hr, the cells were fixed with acetone
and HCV NS5b protein was detected by indirect
immunofluorescence staining using this antibody.

For research use only. Not for clinical diagnosis.

Manufactured by BioAcademia Inc.



COSMO BIO CO., LTD.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

URL: http://www.cosmobio.co.jp e-mail: export@cosmobio.co.jp

[Outside Japan] Phone: +81-3-5632-9617 [国内連絡先] Phone: +81-3-5632-9610 FAX: +81-3-5632-9619