

Anti-POGZ antibody, rabbit polyclonal

BACKGROUND

POGZ (pogo transposable element-derived protein with zinc finger domain; 1,410 aa, 155 kDa) plays a role in mitotic cell cycle progression and is involved in kinetochore assembly and mitotic sister chromatid cohesion. Probably through its association with CBX5 (HP1 alpha) it plays a role in mitotic chromosome segregation by regulating aurora kinase B/AURKB activation and AURKB and CBX5 dissociation from chromosome arms.

Applications

1. Western Blotting (1/1,000 dilution)
2. Immunofluorescence staining (1/500 dilution)
3. Immunoprecipitation (assay dependent)

Not tested for other applications

Immunogen; Recombinant protein expressing 1-562 amino acids of human POGZ.

Reactivity; Reacts specifically with human and mouse POGZ. Not tested in other species

Form; Rabbit antiserum. No azide added. Sterilized by filtration through 0.22 µm membrane.

Size: 100 ul

Storage; Sent at 4°C or -20°C. Store at -20°C. Avoid repeated freeze-thaw cycle.

Data links; [Entrez Gene: 23126](#) Human [SwissProt: Q7Z3K3](#) Human

Reference; This antibody was used in the following article.

Nozawa R.S et al (2010) Human POGZ modulates dissociation of HP1alpha from mitotic chromosome arms through Aurora B activation. Nat. Cell Biol. 12:719-727

[PubMed: 20562864](#)

Anti-POGZ antibody, rabbit polyclonal




Fig.1 Western blot of POGZ

Lane 1: Crude extract of HeLa cells transfected with control siRNA

Lane 2: Crude extract of HeLa cells transfected with POGZ specific siRNA

The antibody was used at 1/1,000 dilution.

Specificity of the antibody to POGZ is confirmed by the disappearance of the POGZ band in the cells treated with the POGZ specific siRNA (lane 2)




Fig.2 Immuno-fluorescence staining of POGZ.

Paraformaldehyde fixed HeLa cells were stained with Hoechst 33342 and anti-POGZ antibody. The antibody was used at 1/500 dilution. Scale bar, 10 μ m




Fig. 3 Immuno-fluorescence staining of POGD on chromosomes.

Chromosome spreads were prepared from HeLa cells and stained with Hoechst 33342 and anti-POGZ antibody. The antibody was used at 1/500 dilution. Scale bar, 10 μ m

For research use only. Not for clinical diagnosis.

Manufactured by BioAcademia, Inc.



COSMO BIO Co., LTD.
Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

http://www.cosmobio.co.jp/index_e.asp

Phone : +81-3-5632-9617

E-mail: export@cosmobio.co.jp

FAX : +81-3-5632-9618