



MONOCLONAL ANTIBODY

For research use only. Not for clinical diagnosis.

Catalog No. BAM-73-505

Anti-ATF6 α antibody, (37-1)

BACKGROUND

ATF6 (activating transcription factor 6) is an endoplasmic reticulum (ER) membrane-bound transcription factor activated in response to ER stress. When unfolded proteins accumulate in the ER, **ATF6** is cleaved by regulated intramembrane proteolysis. The resulting amino-terminal fragment translocates to the nucleus and activates transcription by binding to ER stress-response elements present in the promoter regions of ER stress-inducible genes including those encoding ER chaperones and components of ER-associated degradation. **ATF6** consists of two closely related factors, **ATF6 α** and ATF6 β , in mammals. **ATF6 α** but not ATF6 β plays a pivotal role in transcriptional control.

Product type	Primary antibodies
Host	Mouse
Source	
Form	Liquid Purified monoclonal antibody (IgG) 1mg/ml in PBS, 50% glycerol, filter-sterilized
Volume	100 μ g
Concentration	
Specificity	ATF6 α
Antigen	Recombinant ATF6 α (amino-terminal fragment of ATF6 fused to GST) Epitope: not determined
Clone	37-1
Isotype	Mouse IgG1 κ

Application notes WB, Immunoprecipitation (IP) (less efficient than clone1-7)
This antibody does not work for immunofluorescence analyses.

Recommended use

Recommended dilutions

Optimal dilutions/concentrations should be determined by the end user.
Data Link: Swiss-Prot [P18850] (human ATF6 alpha)

Cross reactivity human and mouse ATF6 α

Storage -20°C (long period, -70°C)

- References**
- 1) Hai T, *et al.*, (1989) "Transcription factor ATF cDNA clones: an extensive family of leucine zipper proteins able to selectively form DNA-binding heterodimers." *Genes Dev* **3**: 2083-2090 [PMID: 2516827]
 - 2) Haze K, *et al.*, (1999) "Mammalian transcription factor ATF6 is synthesized as a transmembrane protein and activated by proteolysis in response to endoplasmic reticulum stress". *Mol Biol Cell* **10**: 3787-3799 [PMID: 10564271]
 - 3) Yamamoto K. *et al.*, (2007) "Transcriptional induction of mammalian ER quality control proteins is mediated by single or combined action of ATF6 and XBP1". *Dev. Cell* **13**: 365-376 [PMID: 17765680]



Protocol for ATF6 α analysis using anti-human ATF6 α monoclonal antibody (37-1)

Both endogenous precursor ATF6 α , pATF6 α (P), and its cleaved product, pATF6 α (N), can be detected in human cells such as HeLa cells by western blot analysis using anti-human ATF6 α monoclonal antibody clone 37-1 (Fig. 1), according to the procedures described below.

As clone 37-1 cross reacts with mouse ATF6 α , both endogenous precursor ATF6 α , pATF6 α (P), and its cleaved product, pATF6 α (N), can be detected in mouse cells such as NIH3T3 cells by western blot analysis (Fig. 2), according to the procedures described below.




Fig.1 Western blot analysis of human cell extracts using this antibody: Conversion of pATF6 α (P) to pATF6 α (N) in DTT- or tunicamycin-treated cells.

- 1) untreated
- 2) DTT: 1mM dithiothreitol (reducing reagent) for 1 h.
- 3) Tm: 2 μ g/ml tunicamycin (inhibitor of N-glycosylation) for 3 h.
- 4) Tm: 2 μ g/ml tunicamycin (inhibitor of N-glycosylation) for 7 h.

The asterisk denotes an unglycosylated form of pATF6 α (P).

ATF6 α is constitutively expressed as pATF6 α (P) (~90-kDa protein), and converted to pATF6 α (N) (>50-kDa protein) in ER-stressed cells.




Fig.2 Western blot analysis of mouse cell extracts using this antibody: Conversion of pATF6 α (P) to pATF6 α (N) in DTT- or tunicamycin-treated cells.

- 1) untreated.
- 2) DTT: 1mM dithiothreitol for 1 h.
- 3) Tm: 2 μ g/ml tunicamycin for 3 h.
- 4) Tm: 2 μ g/ml tunicamycin for 7 h.

The asterisk denotes an unglycosylated form of pATF6 α (P).

ATF6 α is constitutively expressed as pATF6 α (P) (~90-kDa protein), and converted to pATF6 α (N) (>50-kDa protein) in ER-stressed cells.



Western blotting

SDS-sample buffer: 50 mM Tris/HCl, pH6.8, containing 2% SDS, (100 mM DTT), 10% glycerol and BPB
PBST: PBS containing 0.1% Tween 20

Blocking buffer: PBS containing 0.1% Tween 20 and 5% skim milk

• Sample Preparation (for HeLa or NIH3T3 cells cultured in 6cm dish)

- (1) Wash cells with ice-cold PBS.
- (2) Scrape cells in 500 μ l of ice-cold PBS (+ protease inhibitor cocktail and 10 μ M MG132) 2 times and collect cells by centrifugation at 5,000 rpm for 2 min.
- (3) Lyse cells directly in 100 μ l of SDS-sample buffer without reducing reagent (+ protease inhibitor cocktail and 10 μ M MG132).
- (4) Vortex mix vigorously.
- (5) Boil the lysate for 5 min and vortex well.
- (6) If the lysate is still viscous, boil again and vortex mix vigorously.
- (7) Centrifuge at 14,000 rpm for 2 min.
- (8) Determine protein concentration using BCA protein assay kit.

• SDS-PAGE and incubation with antibody

- (9) Add one-tenth volume of 1 M DTT and boil for 5 min.
- (10) Subject 50 μ g of the lysate to 8% SDS-PAGE.
- (11) Transfer to nitrocellulose membrane (such as Hybond-ECL, GE Healthcare).
- (12) Incubate the membrane in Blocking buffer overnight at 4°C. (**overnight incubation is essential**)
- (13) Incubate the membrane with primary antibody diluted in Blocking buffer (1:500-1:1000) for 1 h at room temperature or overnight at 4°C. Wash the membrane 3 times each for 5 min with PBST.
- (14) Incubate the membrane with HRP-conjugated secondary antibody for 1 h at room temperature. We recommend "ECL anti-mouse IgG, Horseradish Peroxidase linked F(ab')₂ fragment" (GE Healthcare NA9310V-1ML).
- (15) Wash the membrane 3 times each for 5 min with PBST.
- (16) Detect signals using an appropriate luminescent reagent.

*Clearer results can be obtained by using 'Can Get Signal (cat# TYB-NKB-101T)' during incubation with primary and secondary antibodies, according to the manufacture's instructions.

For research use only. Not for clinical diagnosis.

Manufactured by BioAcademia, Inc.



COSMO BIO Co., LTD.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

URL: <http://www.cosmobio.co.jp>

e-mail: export@cosmobio.co.jp

[Outside Japan] Phone : +81-3-5632-9617

[国内連絡先] Phone : +81-3-5632-9610

FAX : +81-3-5632-9618

FAX : +81-3-5632-9619