



**MONOCLONAL ANTIBODY**

*For research use only. Not for clinical diagnosis.*

**Catalog No. AGE-M02**

# Anti N<sup>ε</sup>-(carboxyethyl) lysine (CEL)

## BACKGROUND

N<sup>ε</sup>-(carboxyethyl) lysine (CEL) is generated from protein modification by methylglyoxal (MG), which is enzymatically derived from the Embden-Meyerhof and polyol pathways, through the degradation of glyceraldehyde-3-phosphate (G3P) (Phillips and Thornalley, 1993). McLellan et al. (McLellan et al., 1994) demonstrated that plasma MG concentrations in insulin-dependent diabetic patients were 7-times higher than those of healthy individuals.

<b>Product type</b>	Primary antibodies
<b>Host</b>	Mouse
<b>Source</b>	Purified from ascite
<b>Form</b>	Liquid with 0.1% proclin
<b>Volume</b>	100 ul
<b>Concentration</b>	0.2 mg/ml
<b>Specificity</b>	CEL
<b>Antigen</b>	CEL-BSA
<b>Clone</b>	CEL-SP
<b>Isotype</b>	IgG1

## Application notes

### Recommended use

WB, IHC, ELISA



### Recommended dilutions

Western blotting, 1/200 to 1/400

Immunohistochemistry, 1/100 to 1/200

ELISA, 1/200 to 1/400


Optimal dilutions/concentrations should be determined by the end user.



**CEL**



CEL  
production  
pathway



Storage

Store below -20°C (below -70°C for prolonged storage).

Aliquot to avoid cycles of freeze/thaw.

References

- 1) Nagai R., Fujiwara Y., Mera K., Yamagata K., Sakashita N., Takeya M. Immunochemical detection of N<sup>ε</sup>-(carboxyethyl)lysine using a specific antibody. J. Immunol. Methods 332, 112-120 (2008)

*For research use only. Not for clinical diagnosis.*



COSMO BIO Co., LTD.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

URL: <http://www.cosmobio.co.jp>

e-mail: [export@cosmobio.co.jp](mailto:export@cosmobio.co.jp)

[Outside Japan] Phone : +81-3-5632-9617

[国内連絡先] Phone : +81-3-5632-9610

FAX : +81-3-5632-9618

FAX : +81-3-5632-9619