

POLYCLONAL ANTIBODY


For research use only. Not for clinical diagnosis.

Catalog No. CNP- WB-229

Anti NAGA

Product type	Primary antibodies
Immunogen	NAGA (NP_000253.1, 1 a.a. ~ 411 a.a) full-length human protein.
Form	Liquid Protein A purified
Volume	100ul
Specificity	Human
Host	Rabbit
Immunogen Sequence	MLLKTVLLLGHVAQVLMIDNGLLQTPPMGWLAWERFRCNINCEDEPKNCISEQLFMEMAD RMAQDQWRDMGYTYLNIDDCWIGGRDASGRLMPDPKRFPHGIPFLADYVHSLGLKLGIIYA DMGNFTCMGYPGTTLDKVVQDAQTFAEWKVDMLKLDGCFSTPEERAQGYPKMAAALNATG RPIAFSCSWPAYEGGLPPRVNYSLLADICNLWRNYDDIQDSWWSVLNWFVEHQDILQ PVAGPGHWNDPDMLLIGNFGLSLEQSRAQMALWTVLAAPLLMSTDLRTISAQNMDILQNP LMIKINQDPLGIQGRRIHKEKSLIEVYMRPLSNKASALVFFSCRTDMPYRYHSSLGQLNF TGSVIYEAQDVYSGDIISGLRDETNTFTVIINPSGVVMWYLYPIKNLEMSQQ
Gene Alias	D22S674, GALB
Application notes	WB Antibody dilution: 1:1000 ~ 1:2000. Specific band of ~46.60 kDa using NAGA transfected 293T lysate.

Western Blotting (Transfected lysate)




Western Blot analysis of NAGA expression in transfected 293T cell line by NAGA rabbit polyclonal antibody.

Lane 1: NAGA transfected lysate(46.60 KDa).
Lane 2: Non-transfected lysate.

Antibody dilution: 1:1000. Western Blot analysis of NAGA expression in human liver (Predicted Mw: 43.7480 kDa).

Western Blotting (Tissue lysate)



NAGA rabbit polyclonal antibody. Western Blot analysis of NAGA expression in human liver.

Storage

For continuous use, store at 2-8°C for one-two days. For extended storage, store in -20°C freezer. Working dilution samples should be discarded if not used within 12 hours.

For research use only. Not for clinical diagnosis.



COSMO BIO Co., LTD.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

URL: <http://www.cosmobio.co.jp>

e-mail: export@cosmobio.co.jp

[Outside Japan] Phone : +81-3-5632-9617

[国内連絡先] Phone : +81-3-5632-9610

FAX : +81-3-5632-9618

FAX : +81-3-5632-9619