

Cosmo Bio Co., Ltd

TOYO EKIMAE BLDG, 2-20,TOYO 2CHOME, KOTO-KU,TOKYO 135-0016,JAPAN

TEL: +81-3-5632-9617 FAX: +81-3-5632-9618

URL: http://www.cosmobio.co.jp/e-mail: export@cosmobio.co.jp

国内連絡先

TEL: 03-5632-9610 FAX: 03-5632-9619

Catalog No.CBX00278

Mouse monoclonal antibody Anti-Human HARS

Formulation

Mouse monoclonal anti-human **HARS** antibody in PBS (3.0 mM KCl, 1.5 mM KH₂PO₄, 140 mM NaCl, 8.0 mM Na₂HPO₄ (pH 7.4)) containing 1% bovine serum albumin (BSA) and 0.05% sodium azide (NaN₃).

■ Antibody concentration

 $100 \mu g/ml (1.0 ml)$

Storage

Store at 2-8°C for up to one year. We recommend storing at −20°C for long-term storage. Avoid repeat freezing and thawing cycles.

Preparation

This antibody was purified using protein G column chromatography from culture supernatant of hybridoma cultured in a medium containing bovine IgG-depleted (approximately 95%) fetal bovine serum.

Sterility

Filtered through a 0.22 µm membrane.


Applications

WB,IP,IC,Dot Blot

Disposal

This antibody solution contains sodium azide (NaN₃) as a preservative. There is a potential hazard that NaN₃ reacts with copper or lead to produce an explosive compound. For safe disposal, the vial has to be washed thoroughly with water.

Lot No. HARSA6-2 Clone No. HARSA6 Antibody class: IgG1



Safety warnings and precautions

Caution must be taken to avoid contact with skin or eyes. In such a case, rinse thoroughly at once with water. Do not ingest, inhale, or swallow. Seek medical attention immediately.

Wear appropriate protective clothing such as laboratory overalls, safety glasses and gloves.

It is strongly advised that this product should be handled by people who have been well trained in laboratory techniques and that it is handled with care pursuant to the principles of good laboratory practice. All chemicals are deemed potentially harmful.

The vial is prone to fall over. Use caution, especially when the lid is off.



Catalog No. CBX00278

Mouse monoclonal antibody Anti-Human HARS


■ Background

Aminoacyl-tRNA synthetases are a class of enzymes that charge tRNAs with their cognate amino acids. The protein encoded by this gene is a cytoplasmic enzyme which belongs to the class II family of aminoacyl-tRNA synthetases. The enzyme is responsible for the synthesis of histidyl-transfer RNA, which is essential for the incorporation of histidine into proteins. The gene is located in a head-to-head orientation with HARSL on chromosome five, where the homologous genes share a bidirectional promoter. The gene product is a frequent target of autoantibodies in the human autoimmune disease polymyositis/dermatomyositis. [NCBI Entrez Gene Summary]

■ Recommended condition

WB: 0.2-2 μg/ml IP: 100-500 μg/sample FC: 0.5-2 μg/sample ICC: 2-100 μg/ml

Application




250 -150 -100 -75 -50 -37 -

kDa


Detection of HARS by Western blot. Samples: Whole cell lysate from human HeLa (H, 25 $\mu g)$, mouse NIH3T3 (M, 25 $\mu g)$ and rat F2408 (R, 25 $\mu g)$ cells. [Lot No. HARSA6-2]

Predicted molecular weight: 57 kDa

Immunoprecipitation: RIPA lysate of HeLa cells was incubated with anti-HARS mAb. [Lot No. HARSA6-1]



HeLa cells were fixed in 2% paraformaldehyde/PBS and then permeabilized in 90% methanol. Cells were stained with anti-HARS mAb (shaded) or isotype control (unshaded) followed by Alexa Fluor® 488-conjugated goat anti-mouse IgG. [Lot No. HARSA6-2]



Immunostaining analysis in HeLa cells. HeLa cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100 in PBS. The cells were immunostained with anti-HARS mAb. [Lot No. HARSA6-1]