

Cosmo Bio Co., Ltd

TOYO EKIMAE BLDG, 2-20,TOYO 2CHOME, KOTO-KU,TOKYO 135-0016,JAPAN

TEL: +81-3-5632-9617 FAX: +81-3-5632-9618

URL: http://www.cosmobio.co.jp/e-mail: export@cosmobio.co.jp

国内連絡先

TEL: 03-5632-9610 FAX: 03-5632-9619

Catalog No.CBX00501

Mouse monoclonal antibody Anti-Human KNTC2

■ Formulation

Mouse monoclonal anti-human **KNTC2** antibody in PBS (3.0 mM KCl, 1.5 mM KH₂PO₄, 140 mM NaCl, 8.0 mM Na₂HPO₄ (pH 7.4)) containing 1% bovine serum albumin (BSA) and 0.05% sodium azide (NaN₃).

Antibody concentration

 $100 \mu g/ml (1.0 ml)$

■Storage

Store at 2-8°C for up to one year.

We recommend storing at -20° C for long-term storage. Avoid repeat freezing and thawing cycles.

Preparation

This antibody was purified using protein G column chromatography from culture supernatant of hybridoma cultured in a medium containing bovine IgG-depleted (approximately 95%) fetal bovine serum.

Sterility

Filtered through a 0.22 µm membrane.

Applications


WB, IC, Dot Blot

Disposal

This antibody solution contains sodium azide (NaN₃) as a preservative. There is a potential hazard that NaN₃ reacts with copper or lead to produce an explosive compound. For safe disposal, the vial has to be washed thoroughly with water.

All chemicals are deemed potentially harmful.

Lot No. 2497C3a-1 Clone No. 2497C3a Antibody class: IgG1



Safety warnings and precautions

Caution must be taken to avoid contact with skin or eyes. In such a case, rinse thoroughly at once with water. Do not ingest, inhale, or swallow. Seek medical attention immediately.

Wear appropriate protective clothing such as laboratory overalls, safety glasses and gloves.

It is strongly advised that this product should be handled by people who have been well trained in laboratory techniques and that it is handled with care pursuant to the principles of good laboratory practice.

The vial is prone to fall over. Use caution, especially when the lid is off.



Catalog No. CBX00501

Mouse monoclonal antibody Anti-Human KNTC2


Background

HEC is one of several proteins involved in spindle checkpoint signaling. This surveillance mechanism assures correct segregation of chromosomes during cell division by detecting unaligned chromosomes and causing prometaphase arrest until the proper bipolar attachment of chromosomes is achieved.[supplied by OMIM] [NCBI Entrez Gene Summary]

Recommended condition


WB: $0.2-2 \mu g/ml$ IC: $2-100 \mu g/ml$

Application



Detection of human KNTC2 by Western blot. Samples: Whole cell lysate (25 $\,\mu$ g) from HeLa cells. [Lot No. 2497C3a-1]

Predicted molecular weight: 73 kDa



Immunostaining analysis in HeLa cells. HeLa cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100 in PBS. The cells were immunostained with anti-KNTC2 mAb. Nuclear was stained with Hoechst (blue fluorescence). [Lot No. 2497C3a-1]