

Anti-Medaka vitellogenin IgG

Horseradish Peroxidase conjugate

| | |
|-------------------------------|--|
| Background: | Vitellogenin, the egg yolk precursor protein, is induced by estrogens. On the other hand, environmental estrogens are known to disrupt endocrine system in animals. Recently, vitellogenin has proven to be an ideal marker for environmental estrogens. |
| Code: | MVD33-HRP |
| Lot: | OH06 |
| lot size: | 0.5 mL |
| Presentation: | Supplied as a liquid in 0.01M Tris , 0.15M NaCl, 1% BSA, pH 7.5 |
| Immunogen Source: | purified Medaka vitellogenin |
| Application: | ELISA, Western blot |
| Normal dilution range: | ELISA: 1:100 - 1:1000 Western blot: 1:100 - 1:5000 |
| Storage: | at -20° C for at least 12 months Avoid freeze thaw cycles |
| Notes: | Since assay conditions vary, the optimum dilution should be determined for particular application. |
| Warning: | For research use only |

COSMO BIO CO.,LTD

TOYO EKIMAE BLDG, 2-20, TOYO 2CHOME, KOTO-KU, TOKYO 135-0016, JAPAN
TEL: +81-3-5632-9605 FAX: +81-3-5632-9614 e-mail: export@cosmobio.co.jp

www.cosmobio.co.jp