

The RI Assay of anti 5alpha-Androstane-3alpha,17beta-diol-17-Succinate-BSA according to FKA-130

The Cross reactivities of anti 5alpha-Androstane-3alpha,17beta-diol-17-Succinate-BSA (FKA-130) by RI Assay

Antigen	5alpha-Androstane-3alpha,17beta-diol-17-Succinate-BSA	
Antibody	FKA-130 Serum	Freeze-drying of 100µl / Vial
Dilution of Serum	2000test for RI Assay	250µl/tube use
Binding of Total	60%	
Teststerone dose of B0/2	300 pg	
Strage/Handling	Maintain at 4°C-8 °C (-20 °C for long term storage) in undiluted aliquots. Avoid repeated freeze/thaw cycles.	

Steroid	Cross Reaction
5alpha-Androstane-3alpha,17beta-diol	100 %
5alpha-Dihydrotestosterone	1.60 %
5alpha-Androstane-3,17-dione	1.48 %
5-Androstene-3beta.17beta-diol	1.20 %
5-Androstene-3.17-dione	0.60 %
Testosterone	0.20 %
Pregnenolone	0.02 %
Progesterone	0.01 %
17alpha-OH-Progesterone	0.01 %
Corticosterone	0.01 %
Cortisol	0.01 %
Estradiol	0.01 %

For research use only; not for use as a diagnostic.

COSMO BIO CO.,LTD

TOYO EKIMAE BLDG, 2-20,TOYO 2CHOME,KOTO-KU,TOKYO 135-0016,JAPAN
 TEL: +81-3-5632-9605 FAX: +81-3-5632-9614 e-mail: export@cosmobio.co.jp

www.cosmobio.co.jp