



Code No.KAL-KH011-02-EX

For research use only

**Advanced Glycation End Products (AGE)
Anti CML Monoclonal Antibody (Clone No. CMS-10)
Peroxidase conjugated**

Reaction of protein amino groups with glucose leads, through the early products such as a Schiff base and Amadori rearrangement products, to the formation of advanced glycation end products (AGE). Recent immunological studies using anti-AGE antibody (6D12) demonstrated the presence of AGE-modified proteins in several human tissues: (i) human lens (nondiabetic and noncataractous), (ii) renal proximal tubules in patients with diabetic nephropathy and chronic renal failure, (iii) diabetic retina, (iv) peripheral nerves of diabetic neuropathy, (v) atherosclerotic lesions of arterial walls, (vi) β_2 -microglobulin forming amyloid fibrils in patients with hemodialysis-related amyloidosis, (vii) senile plaques of patients with Alzheimer's disease, (viii) the peritoneum of CAPD patients, (ix) skin elastin in actinic elastosis, and (x) ceroid/lipofuscin deposits. These results suggest a potential role of AGE-modification in normal aging as well as age-enhanced disease processes. This antibody named as 6D12 has been used to demonstrate AGE-modified proteins in these human tissues, indicating potential usefulness of this antibody for histochemical identification and biochemical quantification of AGE-modified proteins.


N^ϵ -(carboxymethyl)lysine (CML) was a major AGE structure identified by Banes et al. in 1989. Oxidative cleavage of Amadori products is considered as a major route to CML formation in vivo. Banes also revealed that CML was directly formed from the reaction between lipidoxidative products and Lysine residue. Thus, CML could become a marker of oxidative stress and long term damage to protein in aging, atherosclerosis, and diabetes.

Package Size	50 μ g (200 μ L/vial)
Format	Mouse monoclonal antibody, Peroxidase conjugated 0.25 mg/mL
Buffer	Block Ace as a stabilizer, containing 0.1% Proclin as a bacteriostat
Storage	Store below -20°C . Once thawed, store at 4°C . Repeated freeze-thaw cycles should be avoided.
Clone No.	CMS-10
Subclass	IgG1
Purification method	The splenic lymphocytes from BALB/c mouse, immunized with CML-KLH were fused to myeloma P3U1 cells. The cell line (CMS-10) with positive reaction was grown in ascitic fluid of BALB/c mouse, from which the antibody was purified by Protein G affinity chromatography.

Working dilution for immunohistochemistry: 5-10 μ g/mL; for ELISA: 0.1-1.0 μ g/mL

N^ϵ — (carboxymethyl) lysine

CML





Code No.KAL-KH011-02-EX

**Advanced Glycation End Products (AGE)
Anti CML Monoclonal Antibody (Clone No. CMS-10)
Peroxidase conjugated**

【References】

1. Dunn JA, Patrick JS, Thorpe SR, Baynes JW (1989): Oxidation of glycated proteins: Age-dependent accumulation of N^ε-(carboxymethyl) lysine in lens proteins. *Biochemistry*. 28: 9464-9468.
2. Fu MX, Requena JR, Jenkins AJ, Lions TJ, Baynes JW, Thorpe SR(1996): The advanced glycation end product, N^ε-(carboxymethyl) lysine, is a product of both lipid peroxidation and glycoxidation reactions. *J.Biol.Chem.*271: 9982-9986

* These references are the background of CML, and are not this antibody examples.

Distributor



COSMO BIO Co., LTD.
Inspiration for Life Science

TOYO EKIMAE BLDG. 2-20, TOYO 2CHOME
KOTO-KU, TOKYO 135-0016, JAPAN
TEL : +81-3-5632-9617
FAX : +81-3-5632-9618
URL : <http://www.cosmobio.co.jp/>
e-mail : export@cosmobio.co.jp

Manufacturer



Trans Genic Inc.

7-1-6 Minatojimaminami-machi,
Chuo-ku, Kobe,650-0047 JAPAN
TEL : +81-78-306-0590
FAX : +81-78-306-0589
URL : <http://www.transgenic.co.jp/>
e-mail : techstaff@transgenic.co.jp