



Code No.KAL-KH025-01-EX

For research use only

Advanced Glycation End Products (AGE)
Anti CEL Monoclonal Antibody (Clone No. KNH-30)
Biotin conjugated

Reaction of protein amino groups with glucose leads, through the early products such as a Schiff base and Amadori rearrangement products, to the formation of advanced glycation end products (AGE). Recent immunological studies using anti-AGE antibody (6D12) demonstrated the presence of AGE-modified proteins in several human tissues: (i) human lens (nondiabetic and noncataractous), (ii) renal proximal tubules in patients with diabetic nephropathy and chronic renal failure, (iii) diabetic retina, (iv) peripheral nerves of diabetic neuropathy, (v) atherosclerotic lesions of arterial walls, (vi) β_2 -microglobulin forming amyloid fibrils in patients with hemodialysis-related amyloidosis, (vii) senile plaques of patients with Alzheimer's disease, (viii) the peritoneum of CAPD patients, (ix) skin elastin in actinic elastosis, and (x) ceroid/lipofuscin deposits. These results suggest a potential role of AGE-modification in normal aging as well as age-enhanced disease processes. This antibody named as 6D12 has been used to demonstrate AGE-modified proteins in these human tissues, indicating potential usefulness of this antibody for histochemical identification and biochemical quantification of AGE-modified proteins.


CEL is known to generate from protein modification by methylglyoxal . Mclellan et al. demonstrated that plasma methylglyoxal, which is believed to be generate from Embden-Meyerhof and polyol pathways, concentrations in insulin-dependent diabetic patients were about 7-times higher than those of normal individuals. For examples, CEL was identified in human lens proteins at a concentration similar to that of CML and its accumulation increased with age like CML, indicating that CEL may play an important marker for aging and age-dependent disease such as diabetic complications.

Package Size	50 μ g (200 μ L/vial)
Format	Mouse monoclonal antibody , Biotin conjugated 0.25 mg/mL
Buffer	Block Ace as a stabilizer, containing 0.1% Proclin as a bacteriostat
Storage	Store below -20°C . Once thawed, store at 4°C . Repeated freeze-thaw cycles should be avoided.
Clone No.	KNH-30
Subclass	IgG1
Purification method	The splenic lymphocytes from BALB/c mouse, immunized with CEL-BSA were fused to myeloma P3U1 cells. The cell line (KNH-30) with positive reaction was grown in ascitic fluid of BALB/c mouse, from which the antibody was purified by Protein G affinity chromatography.

Working dilution for immunohistochemistry: 5-10 μ g/mL; for ELISA: 0.1-1.0 μ g/mL

N^ε— (carboxyethyl) lysine

CEL





Code No.KAL-KH025-01-EX

Advanced Glycation End Products (AGE)
Anti CEL Monoclonal Antibody (Clone No. KNH-30)
Biotin conjugated

【References】

1. Ahmed MU, Brinkmann E, Degenhardt TP, Thorpe SR, Baynes JW: N^ε-(Carboxyethyl)lysine, a product of the chemical modification of proteins by methylglyoxal, increases with age in human lens proteins. *Biochem J* 324:565-570, 1997
2. Degenhardt TP, Thorpe SR, Baynes JW: Chemical modification of proteins by methylglyoxal. *Cell Mol Biol* 44:1139-1145, 1998
3. Mclellan AC, Thornalley PJ, Benn J, Sonksen PH: Glyoxalase system in clinical diabetes mellitus and correlation with diabetic complications. *Clinical Science* 87: 21-29, 1994

*These references are the background of CEL, and are not this antibody examples

Distributor



COSMO BIO Co., LTD.
Inspiration for Life Science

TOYO EKIMAE BLDG. 2-20, TOYO 2CHOME
KOTO-KU, TOKYO 135-0016, JAPAN
TEL : +81-3-5632-9617
FAX : +81-3-5632-9618
URL : <http://www.cosmobio.co.jp/>
e-mail : export@cosmobio.co.jp

Manufacturer



Trans Genic Inc.

7-1-6 Minatojimaminami-machi,
Chuo-ku, Kobe,650-0047 JAPAN
TEL : +81-78-306-0590
FAX : +81-78-306-0589
URL : <http://www.transgenic.co.jp/>
e-mail : techstaff@transgenic.co.jp