



## TRAP Staining Kit

### Introduction

The TRAP Staining Kit (Cat.No.PMC-AK04F-COS) is used for the staining of Tartrate-Resistant Acid Phosphatase in osteoclasts. Bone mass is controlled by the balance between the activity of osteoblasts bone formation and the activity of osteoclasts. Alkaline phosphatase (ALP: ALP staining kit, Cat.No.PMC-AK20-COS) and tartrate-resistant acid phosphate are used as markers for osteoblasts and osteoclasts, respectively.

### Components

| Component                  | Quantity | Storage |
|----------------------------|----------|---------|
| Tartrate-containing Buffer | 50mL     | 4°C     |
| Chromogenic Substrate      | 10vials  | 4°C     |

**One kit contains reagents for 10 × 96-well plates**

### Additional Materials Required

- 10% Formalin Neutral Buffer Solution
- Phosphate Buffered Saline (PBS)
- Distilled or deionized water

### Protocol (96-well plate format)

1. Remove culture medium. Wash each well once with 100uL of PBS.
2. Add 50uL of the 10% Formalin Neutral Buffer Solution to each well and fix for 5 minutes at room temperature.
3. Wash each well with 250uL of deionized water. (× 3 times)
4. Dissolve 1 vial of Chromogenic Substrate with 5mL of Tartrate-containing Buffer.
5. Add 50uL of Chromogenic Substrates to each well.
6. Incubate at 37°C for 20-60 minutes. Adjust incubation time until stained TRAP is clearly showing the result in figure 1.
7. Wash with deionized water to stop the reaction.

**Note:** Excess incubation will be cause of over stained.




Figure 1: TRAP staining of Osteoclast



COSMO BIO Co., LTD.

Inspiration for Life Science

TRAP Staining Kit Catalog No. PMC-AK04F-COS

## References

- (1) Sunao Takesita, Keisuke Kaji, Akira Kudo. Identification and Characterization of the New Osteoclast Progenitor with Macrophage Phenotypes Being Able to Differentiate into Mature Osteoclasts. JOURNAL OF BONE AND MINERAL RESEARCH Volume 15, Number 8 (2000). 1477-1488.
- (2) BEN A. A. SCHEVEN, JONE S. MILNE, SIMON P. ROBINS. A SEQUENTIAL CULTURE APPROACH TO STUDY OSTEOCLAST DIFFERENTIATION FROM NONADHERENT PORCINE BONE MARROW CELLS. In Vitro Cell. Dev. Biol. July-August (1998). Animal 34: 568-577.
- (3) Martha J. Somerman, Janice E. Berry, Zhila Khalkhali-Ellis, Philip Osdoby, Robert U. Simpson. Enhanced Expression of  $\alpha_v$  Integrin Subunit and Osteopontin during Differentiation of HL-60 Cells along Monocytic Pathway. EXPERIMENTAL CELL RESEARCH 216, 335-341(1995).
- (4) Ichiro Itonaga, Afsie Sabokbar, Susan D. Neale, Nicholas A. Athanasou. 1,25-Dihydroxyvitamin D<sub>3</sub> and Prostaglandin E<sub>2</sub> Act Directly on Circulating Human Osteoclast Precursors. Biochemical and Biophysical Research Communications 264, 590-595(1999).
- (5) Hiroshi Takayanagi, Kouetsu Ogasawara, Shigeaki Hida, Tomoki Chiba, Shigeo Murata, Kojiro Sato, Akinori Takaoka, Taeko Yokochi, Hiromi Oda, Keiji Tanaka, Kozo Nakamura and Tadatsugu Taniguchi. T-cell-mediated regulation of osteoclastogenesis by signaling cross-talk between RANKL and IFN- $\gamma$ . Nature, Vol. 408, No.6812, 600-605, 30 November 2000.
- (6) Morinobu, A., Biao, W., Tanaka, S., Horiuchi, M., Jun, L., Tsuji, G., Sakai, Y., Kurosaka, M., Kumagai, S. (-)-Epigallocatechin-3-Gallate Suppresses Osteoclast Differentiation and Ameliorates Experimental Arthritis in Mice. Arthritis Rheum. 58, 2012-2018 (2008)
- (7) Hase, H., Kanno, Y., Kojima, H., Sakurai, D., Kobata, T. Coculture of Osteoclast Precursors With Rheumatoid Synovial Fibroblasts Induces Osteoclastogenesis via Transforming Growth Factor beta-Mediated Down-Regulation of Osteoprotegerin. Arthritis Rheum. 58, 3356-3365 (2008)
- (8) Cao, H., Kuboyama, N. A Biodegradable Porous Composite Scaffold of PGA/beta-TCP for Bone Tissue Engineering. Bone. 46, 386-395 (2010)
- (9) Okada, Y., Hamada, N., Kim, Y., Takahashi, Y., Sasaguri, K., Ozono, S., Sato, S. Blockade of Sympathetic beta-receptors Inhibits Porphyromonas Gingivalis-induced Alveolar Bone Loss in an Experimental Rat Periodontitis Model. Arch. Oral Biol. 55, 502-508 (2010)
- (10) Okamoto, A., Ohnishi, T., Bandow, K., Kakimoto, K., Chiba, N., Maeda, A., Fukunaga, T., Miyawaki, S., Matsuguchi, T. Reduction of Orthodontic Tooth Movement by Experimentally Induced Periodontal Inflammation in Mice. Eur. J. Oral Sci. 117, 238-247 (2009)

## Notice to Purchaser

This product is to be used for Research Purposes Only. It is not to be used for Drug or Diagnostic Purposes, nor is it intended for Human Use. Primary cell products may not be resold, modified for resale, or used to manufacture commercial products without the express written consent of Primary Cell Co., Ltd.

Except as otherwise expressly set forth in this use manual, Primary Cell does not make any representation or warranties or conditions of any kind, either express or implied, with respect to the products, or information disclosed hereunder, including, but not limited to, the implied warranties of merchantability, fit for a particular purpose, or noninfringement of the intellectual property rights of third parties.

*For research use only. Not for clinical diagnosis.*

Manufactured by **Primary Cell Co., Ltd.**



COSMO BIO Co., LTD.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

URL: <http://www.cosmobio.com>

e-mail: [export@cosmobio.co.jp](mailto:export@cosmobio.co.jp)

[Outside Japan] Phone : +81-3-5632-9617

[国内連絡先] Phone : +81-3-5632-9610

FAX : +81-3-5632-9618

FAX : +81-3-5632-9619