

Anti human PPAR gamma common mouse monoclonal antibody

PPAR gamma: Peroxisome Proliferator-Activated Receptor gamma

Code No	PP-K8713-00	Application / Recommended Concentration	In order to obtain the best results, optimal working dilutions should be determined by each individual user.
Clone No.	K8713	Western Blot	2 ug/mL
Lot.	***	Non reducing Western Blot	Not yet tested
Concentration	1 mg/mL	ELISA	0.1 ug/mL
Volume	100 uL	Immunoprecipitation	Decide by use
Ig Class	G2a	Supershift Assay	100 ug/mL
Description	<p>Peroxisome proliferator-activated receptor gamma (PPARγ; NR1C3) is a member of orphan nuclear receptor. Oxidized metabolites of linoleic acid, 9-hydroxyoctadecenoic acid (9-HODE) and 13-HODE are activators and ligands of PPARγ. PPARγ is expressed in white adipose tissue, intestinal mucosa, colon, spleen, monocytes, macrophages, retina, cartilage, osteoclast and skeletal muscle. PPARγ plays important roles in lipid and glucose metabolism, and have been implicated in obesity-related metabolic diseases such as hyperlipidemia, insulin resistance, and coronary artery disease. Three members were called PPARα, β, γ. Three N-terminal isoforms, called γ1, γ2 and γ3, are known to arise by alternative splicing and promoter usage from the PPARγ gene. RXR is an obligate partner for PPAR.</p>		
Nomenclature	NR1C3	Chromatin immunoprecipitation	Decide by use
Genbank	U79012	Immunohistochemistry	Not yet tested
Origin	<p>Produced in BALB/c mouse ascites after inoculation with hybridoma of mouse myeloma cells (NS-1) and spleen cells derived from a BALB/c mouse immunized with Baculovirus-expressed recombinant human PPAR gamma2 (2-136 aa) .</p>		
Specificity	<p>This antibody specifically recognizes human PPAR gamma1 and 2, and cross reacts with mouse PPAR gamma1 and 2. This antibody does not recognize human PPAR alpha and delta. Not yet tested in other species.</p>		
Purification	Ammonium sulfate fractionation	Storage	Store at 2 - 8 °C up to one month. For long-term storage, the solution may be frozen in working aliquots. Repeated freezing and thawing is not recommended. Storage in a frost-free freezer is not recommended.
Formulation	Physiological saline with 0.1% NaN ₃ as a preservative.	Reference	Tanaka T, <i>et al.</i> J Atheroscler Thromb, 9(5): 233-241, 2002.
		Notes	Sodium azide may react with lead and copper plumbing to form explosive metal azides. Flush with large amounts of water during disposal.

FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.

Not for Diagnostic or Therapeutic use. Purchase of this product does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written consent of Perseus Proteomics Inc. is prohibited.

MADE IN JAPAN

Aug 23, 2006

Anti human PPAR γ common mouse monoclonal antibody

PPAR γ : Peroxisome Proliferator-Activated Receptor

製品コード PP-K8713-00

Clone No. K8713

Lot. A-2

濃度 1mg/mL

容量 100 μ L

Ig class G2a

Nomenclature NR1C3

Genebank U79012

由来 ヒトPPAR γ 2 (2-136 aa) の Baculovirus 発現物を免疫した BALB/c マウスの脾臓細胞と、マウスミエローマ細胞 (NS-1) を融合して得たハイブリドーマを、BALB/c マウスに接種して得られた腹水。

特異性 ヒト PPAR γ 1, 2 と特異的に反応する。ヒト PPAR α , δ には反応しない。マウスPPAR γ 1, 2と交差反応する。その他の動物種との交差反応は未検討。

精製法 硫酸塩析法

溶媒 生理的食塩水(防腐剤として0.1% NaN₃添加)

FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.

本製品は研究目的のみで使用する事ができます。人やそのほか動物の疾病診断、治療・予防に使用することはできません。
研究目的以外で使用する場合は、あらかじめ弊社にご相談ください。

Application 使用濃度は実験にあわせて至適化が必要です。

Western Blot 可
参考使用濃度 2 μ g/mL

非還元 Western Blot 未検討
参考使用濃度 -

ELISA 可
参考使用濃度 0.1 μ g/mL

免疫沈降 可
参考使用濃度 適宜調製してください

Supershift Assay 可
参考使用濃度 100 μ g/mL

クロマチン免疫沈降 可
参考使用濃度 適宜調製してください

免疫染色 未検討
参考使用濃度 -

保存方法 1ヶ月程度の保存の場合は、2~8℃で保存可能です。長期保存の場合は、抗体を小分けした上で、-20℃以下での保存をお勧めします。また、凍結融解を繰り返すと、抗体が劣化し、本来の性能が得られない場合があるため、お避けください。

参考文献 Tanaka T, *et al.* J Atheroscler Thromb, 9(5): 233-241, 2002.

備考 溶媒に含まれるNaN₃は、鉛や銅と反応し爆発性化合物を形成する恐れがあります。廃棄の際には大量の水と一緒に希釈して廃棄してください。

MADE IN JAPAN

Aug 23, 2006