

Anti human PPAR delta mouse monoclonal antibody

PPAR delta: Peroxisome Proliferator-Activated Receptor delta

Code No	PP-K9436-00
Clone No.	K9436
Lot.	***
Concentration	1 mg/mL
Volume	100 uL
Ig Class	G2a
Description	Peroxisome proliferator-activated receptor beta (PPARb, NUC1, PPARd; NR1C2) is a member of orphan nuclear receptor. PPARb is expressed in uterus and placenta. It plays important roles in lipid and glucose metabolism, and have been implicated in obesity-related metabolic diseases such as hyperlipidemia, insulin resistance, and coronary artery disease. Three members were called PPARa, b, g. RXR is an obligate partner for PPAR.

Nomenclature	NR1C2
Genbank	L07592
Origin	Produced in BALB/c mouse ascites after inoculation with hybridoma of mouse myeloma cells (NS-1) and spleen cells derived from a BALB/c mouse immunized with Baculovirus-expressed recombinant human PPAR delta (1-60 aa).
Specificity	This antibody specifically recognizes human PPAR delta but does not recognize human PPAR alpha and gamma. Not yet tested in other species.

Purification	Ammonium sulfate fractionation
Formulation	Physiological saline with 0.1% NaN3 as a preservative.

Application / Recommended Concentration

In order to obtain the best results, optimal working dilutions should be determined by each individual user.

Western Blot	1 ug/mL
Non reducing Western Blot	Not yet tested
ELISA	0.5 ug/mL
Immunoprecipitation	Not yet tested
Supershift Assay	100 ug/mL
Chromatin immunoprecipitation	Not yet tested
Immunohistochemistry	Not yet tested

Storage	Store at 2 - 8 °C up to one month. For long-term storage, the solution may be frozen in working aliquots. Repeated freezing and thawing is not recommended. Storage in a frost-free freezer is not recommended.
---------	---

Reference

Notes	Sodium azide may react with lead and copper plumbing to form explosive metal azides. Flush with large amounts of water during disposal.
-------	---

FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.

Not for Diagnostic or Therapeutic use. Purchase of this product does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written consent of Perseus Proteomics Inc. is prohibited.

MADE IN JAPAN

Aug 23, 2006

Anti human PPAR δ mouse monoclonal antibody

PPAR δ : Peroxisome Proliferator-Activated Receptor δ

製品コード PP-K9436-00

Clone No. K9436

Lot. A-2

濃度 1mg/mL

容量 100 μ L

Ig class G2a

Nomenclature NR1C2

Genebank L07592

由来 ヒトPPAR δ (1-60 aa) の Baculovirus 発現物を免疫した BALB/c マウスの脾臓細胞と、マウスミエローマ細胞 (NS-1) を融合して得たハイブリドーマを、BALB/c マウスに接種して得られた腹水。

特異性 ヒト PPAR δ と特異的に反応する。ヒト PPAR α , γ には反応しない。その他の動物種との交差反応は未検討。

精製法 硫酸塩析法

溶媒 生理的食塩水 (防腐剤として 0.1% NaN₃ 添加)

FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.

本製品は研究目的のみで使用する事ができます。人やそのほか動物の疾病診断、治療・予防に使用することはできません。
研究目的以外で使用する場合は、あらかじめ弊社にご相談ください。

Application 使用濃度は実験にあわせて至適化が必要です。

Western Blot 可
参考使用濃度 1 μ g/mL

非還元 Western Blot 未検討
参考使用濃度 -

ELISA 可
参考使用濃度 0.5 μ g/mL

免疫沈降 未検討
参考使用濃度 -

Supershift Assay 可
参考使用濃度 100 μ g/mL

クロマチン免疫沈降 未検討
参考使用濃度 -

免疫染色 未検討
参考使用濃度 -

保存方法 1ヶ月程度の保存の場合は、2~8 $^{\circ}$ Cで保存可能です。長期保存の場合は、抗体を小分けした上で、-20 $^{\circ}$ C 以下での保存をお勧めします。また、凍結融解を繰り返すと、抗体が劣化し、本来の性能が得られない場合があるため、お避けください。

参考文献

備考 溶媒に含まれるNaN₃は、鉛や銅と反応し爆発性化合物を形成する恐れがあります。廃棄の際には大量の水と一緒に希釈して廃棄してください。

MADE IN JAPAN

Aug 23, 2006