



Anti PACAP38 (Human) Serum

Cat. No. YII-Y040-EX Lot No. 083171013

Description: This antiserum was raised in a rabbit by immunization with a carrier free synthetic PACAP38 (human) peptide. The product vial contains 50 μ L of the titled antiserum obtained by lyophilizing its 0.001 M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with PACAP.

Immunogen: Synthetic PACAP38 (human), carrier free **Host:** Rabbit

Amino Acid Sequence of PACAP38 (human)¹⁾:

HSDGIFTDSY SRYRKQMAVK KYLAAVLGKR YKQRVKNK-NH₂

Product Form: Lyophilized unpurified serum **Size:** 50 μ L

Reconstitution: Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN₃ 0.1%).

Storage: The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing-thawing should be avoided.

Suggested Working Dilution Range: 1:1,000-2,000 (final dilution ~1:14,000) for radioimmunoassay;

1: 200-1,000 for immunohistochemistry (frozen or paraffin sections)²⁾. Optimal dilution should be determined by each laboratory for each application.

Specificity (based on radioimmunoassay): PACAP38 (human) 100%, PACAP27 (human) 0%, VIP 0%, PHI (rat) 0%, PHI (human) 0%, secretin 0%, helodermin 0%, CCK-8 0%, met-enkephalin 0%, leu-enkephalin 0%, dynorphin 0%, oxytocin 0%, vasopressin 0%

Positive Control (immunohistochemistry): Rat colon

Species Tested: Rat, ginseng radix³⁾

REFERENCES:

1) A. Miyata, A. Arimura et al., Isolation of a novel 38 residue-hypothalamic polypeptide which stimulates adenylate cyclase in pituitary cells. *Biochemistry Biophysics Research Communications* 164:567-574, 1989

2) S. Kimura, Y. Ohshige et al., Localization of pituitary adenylate cyclase-activating polypeptide (PACAP) in the hypothalamus-pituitary system in rats: light and electron microscopic immunocytochemical studies. *Journal of Neuroendocrinology* 6: 503-507, 1994

3) T. Ohishi, N. Yanaihara et al., Immunoreactivities of neuropeptides in plants (*Ginseng Radix* and *Panacis Japonici Rhizoma*). *Proceedings of 18th Gut Hormone Conference, Japan Society of Gut Hormones* (Ed) 13:285-291, 1995

FOR RESEARCH LABORATORY USE ONLY

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM

Distributor



COSMO BIO CO., LTD.
Inspiration for Life Science

TOYO EKIMAE BLDG. 2-20, TOYO 2CHOME
KOTO-KU, TOKYO 135-0016, JAPAN
TEL : +81-3-5632-9617
FAX : +81-3-5632-9618
URL : <http://www.cosmobio.co.jp/>
e-mail : export@cosmobio.co.jp

Manufacturer



2480-1 Awakura, Fujinomiya-shi,
Shizuoka 418-0011, JAPAN
Phone: +81-544-22-2771
Fax: +81-544-22-2770
URL : www.yanaihara.co.jp
E-mail : ask@yanaihara.co.jp