



## **Anti EGF (Human) Serum**

**Cat. No. YII-Y230-EX** **Lot No. 10271201**

**Description:** This antiserum was raised in a rabbit by immunization with a carrier free recombinant EGF (human). The product vial contains 50  $\mu$ L of the titled antiserum obtained by lyophilizing its 0.001 M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with EGF (human).

**Immunogen:** Recombinant EGF (human), carrier free **Host:** Rabbit

### **Amino Acid Sequence of EGF (human)<sup>1)</sup>:**

NSDSECPLSH DGYCLHDGVC MYIEALDKYA CNCVVG YIGE RCQYRDLKWW ELR

**Product Form:** Lyophilized unpurified serum **Size:** 50  $\mu$ L

**Reconstitution:** Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN<sub>3</sub> 0.1%).

**Storage:** The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing-thawing should be avoided.

**Suggested Working Dilution Range:** 1:5,000-12,000 (final dilution ~1:84,000) for radioimmunoassay;

1: 1,000-4,000 for immunohistochemistry (frozen or paraffin sections). Optimal dilution should be determined by each laboratory for each application.

**Specificity** (based on radioimmunoassay)<sup>2,3)</sup>: EGF (human) 100%, EGF (rat) 0.006%, EGF (mouse) 0%, TGF- $\alpha$ (human) 0%, IGF-1 (human) 0%, insulin (porcine) 0%, glucagon 0%, ACTH (human) 0%, PHI (human) 0%

**Positive Control** (immunohistochemistry): Human or rat submaxillary.

**Species Tested:** Human, rat

### **REFERENCES:**

- 1) G.I. Bell, N.M. Fong et al., Human epidermal growth factor precursor: cDNA sequence, expression in vitro and gene organization. Nucleic Acids Research. 14:8427-8432, 1986
- 2) S. Oguchi, N. Yanaihara et al., EGF-like immunoreactivity in saliva and gastric juice of children. Proceedings of 9th Gut Hormone Conference, Japan Society of Gut Hormones (Ed.) 1987, 7:383-389, 1987
- 3) S. Oguchi, N. Yanaihara et al., EGF-like immunoreactivity in urine of children. Proceedings of 10th Gut Hormone Conference, Japan Society of Gut Hormones (Ed.) 8:648-653, 1988

### **FOR RESEARCH LABORATORY USE ONLY**

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM

