



Anti Urocortin (3-40) (Mouse, Rat) Serum

Cat. No. YII-Y360-EX Lot No. 90110122

Description: This antiserum was raised in a rabbit by immunization with a carrier free synthetic urocortin (3-40) (mouse, rat) peptide. The product vial contains 50 µL of the titled antiserum obtained by lyophilizing its 0.001 M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with urocortin (mouse, rat).

Immunogen: Synthetic urocortin (3-40) (mouse, rat), carrier free **Host:** Rabbit

Amino Acid Sequence of urocortin (3-40) (mouse, rat)¹⁾:

DDPPLSIDLT FLLLRTLLEL ARTQSQRERA EQNRIIFDSV-NH₂

Product Form: Lyophilized unpurified serum **Size:** 50 µL

Reconstitution: Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN₃ 0.1%).

Storage: The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstituted, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing-thawing should be avoided.

Suggested Working Dilution Range: 1:1,500 (final dilution ~1:10,500) for radioimmunoassay²⁾; 1:1,000-5,000 for immunohistochemistry (frozen and paraffin section)^{3,4,5)}. Optimal dilution should be determined by each laboratory for each application.

Specificity (based on radioimmunoassay): Urocortin (3-40) (mouse, rat) 100%, urocortin (mouse, rat) 164.7%, urocortin (human) ~100%, urocortin (30-40) (mouse, rat) 100%, urocortin (7-40) (mouse, rat) 260%, urocortin (1-22) (mouse, rat) < 0.1%, CRF (human, mouse, rat) 0%, urotensin I (carp) < 0.1%

Positive Control (immunohistochemistry): Rat pituitary gland

Species Tested: Rat

REFERENCES:

- 1) J.Vaughan, C.Donaldson et al., Urocortin, a mammalian neuropeptide related to fish urotensin I and to corticotropin-releasing factor. *Nature*, 378: 287-292, 1995
- 2) K. Iguchi, N Yanaihara et al., Urocortin, its synthesis, antibody production, and biological activity. *ACTH Related Peptides*, 8:53-59, 1997
- 3) Y. Hara, Y. Ueta et al., Increase of urocortin-like immunoreactivity in the rat hypothalamo-neurohypophysial system after salt loading and hypophysectomy. *Neuroscience Letters* 227: 127-130, 1997
- 4) H. Yamamoto, T. Maeda et al., Urocortin-like immunoreactivity in the substantia nigra, ventral tegmental area and Edinger-Westphal nucleus of rat. *Neuroscience Letters* 243: 21-24, 1998
- 5) T. Nishikimi, A. Miyata et al., Urocortin, a member of the corticotropin-releasing factor family, in normal and diseased heart. *American Journal of Physiology • Heart and Circulatory Physiology* 279: H3031-3039, 2000

FOR RESEARCH LABORATORY USE ONLY

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM

Distributor



COSMO BIO CO., LTD.
Inspiration for Life Science

TOYO EKIMAE BLDG. 2-20, TOYO 2CHOME
KOTO-KU, TOKYO 135-0016, JAPAN
TEL : +81-3-5632-9617
FAX : +81-3-5632-9618
URL : <http://www.cosmobio.co.jp/>
e-mail : export@cosmobio.co.jp

Manufacturer



2480-1 Awakura, Fujinomiya-shi,
Shizuoka 418-0011, JAPAN
Phone: +81-544-22-2771
Fax: +81-544-22-2770
URL : www.yanaihara.co.jp
E-mail : ask@yanaihara.co.jp