



Anti IL-1 β (Rat) Serum **Cat. No. YII-YC020-EX**

Description: Interleukin 1 (IL-1) consists of two proteins, IL-1 α and IL-1 β , which are the products of distinct genes, but which recognize the same cell surface receptors. IL-1 α and IL-1 β show approximately 25% identity in the amino acid sequence.¹⁾ Both IL-1 α and IL-1 β are synthesized as 31K Da precursors that are subsequently cleaved into proteins with molecular weights of approximately 17KDa.²⁾³⁾ Among various species, the amino acid sequences of mature IL-1 α is conserved 60% to 70%, while that of the mature IL-1 β conserved 75% to 78%.⁴⁾ Both IL-1 α and IL-1 β have glycosylation sites, but non-glycosylated recombinant products have biological activities similar to the naturally occurring forms of the molecules.⁵⁾⁶⁾

Immunogen: Recombinant IL-1 β (Rat) **Host:** Rabbit

for ELISA: Approximately 0.14-0.4 ng/well of recombinant rat IL-1 α can be detected using an antiserum concentration of dilution ratio x1000.

for WB: An antiserum concentration of dilution ratio x1000 will allow visualization of 0.01-0.003 ng/lane of recombinant rat IL-1 β under reducing condition.

Specificity:

IL-1 β (Rat)	100%,	IL-1 β (Human)	24.8%
IL-1 α (Rat)	<0.1%	IL-1 α (Human)	<0.1%
TNF- α (Rat)	<0.1%	TNF- α (Human)	<0.1%
Other Cytokines	<0.1%		

Related Antisera: Rabbit Anti IL-1 α (Rat) Serum YII-YC010-EX
Goat Anti IL-1 α (Rat) Serum YII-YC011-EX
Goat Anti IL-1 β (Rat) Serum YII-YC021-EX
Anti IL-1 β (Rat) Monoclonal Antibody YII-YC022-EX

Related Peptides: IL-1 β , TNF- α , TNF- β , Other Cytokines

Storage: Keep frozen below -20°C
Avoid repeated freezing-thawing.

REFERENCES:

- 1) Oppenheim JJ, et al., Immunol. Today, 7:45, 1986
- 2) Giri JG, Lomedico PT, Mizel SB., J. Immunol. 134:343-349, 1985
- 3) DJ Hazuda, J Stricker, F Kueppers, PL Simon, and PR Young., J.Biol.Chem., 265:6318-6322, 1990
- 4) Dodds RA, Merry K, Littlewood A, Gowen M.J., Histochem. Cytochem., 42:733-744, 1994
- 5) Dinarello CA., Blood, 77:1627-1652, 1991
- 6) Casagli MC, Borri MG, Bigio M, Rossi R, Nucci D, Bossu P, Boraschi D, Antoni G. Biochem Biophys Res Commun., 162:357-363, 1989

FOR RESEARCH LABORATORY USE ONLY

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM

Distributor



COSMO BIO CO., LTD.
Inspiration for Life Science

TOYO EKIMAE BLDG. 2-20, TOYO 2CHOME
KOTO-KU, TOKYO 135-0016, JAPAN
TEL : +81-3-5632-9617
FAX : +81-3-5632-9618
URL : <http://www.cosmobio.co.jp/>
e-mail : export@cosmobio.co.jp

Manufacturer



2480-1 Awakura, Fujinomiya-shi,
Shizuoka 418-0011, JAPAN
Phone: +81-544-22-2771
Fax: +81-544-22-2770
URL : www.yanaihara.co.jp
E-mail : ask@yanaihara.co.jp