



Testosterone-3(E)-CMO-BSA polyclonal antibody

Product No. FKA-102

The Cross reactivities of anti Testosterone(T)-3E-CMO-BSA (FKA-102) by RI Assay

Antigen	Testosterone-3E-CMO-BSA	
Antibody[FKA-102]	lyophilized Serum	5 μ l / Vial
Dilution of Serum	2000test for RI Assay	250 μ l / tube use
Binding of Total	60-80%	
Testosterone dose of B0/2	40 pg	
Strage/Handling	Maintain at 4°C-8 °C (-20 °C for long term storage) in undiluted aliquots. Avoid repeated freeze/thaw cycles.	

Steroid	Cross Reaction
Testosterone	100 %
5Alpha-Dihydrotestosterone	8.3 %
4-Androstenedione	3.1 %
Androsterone	0.3 %
5-Androstene-3Beta.17Beta-diol	0.15 %
5Alpha-Androstene-3Alpha.17Beta-diol	0.10 %
5Beta-Androstene-3Alpha.17Beta-diol	0.09 %
Cortisol	0.02 %
Corticosterone	0.01 %
Progesterone	0.01 %
Pregnenolone	<0.01 %
17Alpha-Hydroxypregnenolone	<0.01 %
Aldosterone	<0.01 %
Dehydroepiandrosterone	<0.01 %
Estradiol	<0.01 %

For research use only; not for use as a diagnostic.

