



Anti Dehydroepiandrosterone-3-Succinate polyclonal antibody

Product No. FKA-110E

The Cross reactivities of anti Dehydroepiandrosterone-3-Succinate (FKA-110E)
by ELISA with 11-Oxo-Testosterone-3-CMO-HRP (FKA-109)

Antigen	Dehydroepiandrosterone-3-Succinate-BSA		
Antibody(FKA-110E)	Purified by Protein A	lyophilized	2μl / Vial
Dilution of IgG	2000test for ELISA	50μl/well use	
Label (FKA-109)	Dehydroepiandrosterone-3-Succinate-HRP		1ml / Vial
	1,000test for ELISA	50μl/well use	
O.D. (492/600nm) of B0	0.5		
Dehydroepiandrosterone dose of B0/2	100 ng/ml		
Strage/Handling	Maintain at 4°C-8 °C (-20 °C for long term storage) in undiluted aliquots. Avoid repeated freeze/thaw cycles.		

Steroid	Cross Reaction
Dehydroepiandrosterone (DHA)	100 %
DHA-Sulfate	37.00 %
4-Androstene-3,17-dione	76.00 %
Androsterone	16.00 %
Testosterone	0.80 %
5-Androstene-3,17-diol	1.50 %
Dihydrotestosterone(DHT)	3.50 %
Pregnenolone	0.02 %
Progesterone	0.01 %
17Alpha-OH-Progesterone	0.01 %
Corticosterone	0.01 %
Cortisol	0.01 %
Estradiol	0.01 %

For research use only; not for use as a diagnostic.

