



Anti 11-Oxo-Testosterone-3-CMO-BSA polyclonal antibody

Product No. FKA-118

The Cross reactivities of anti 11-Oxo-Testosterone-3-CMO-BSA (FKA-118) by RI Assay

Antigen	11-Oxo-Testosterone-3-CMO-BSA	
Antibody[FKA-118]	lyophilized Serum	20 μ l / Vial
Dilution of Serum	2000test for RI Assay	250 μ l / tube use
Binding of Total	51%	
Testosterone dose of B0/2	200 pg	
Strage/Handling	Maintain at 4°C-8 °C (-20 °C for long term storage) in undiluted aliquots. Avoid repeated freeze/thaw cycles.	

Steroid	Cross Reaction
11-Oxo-Testosterone	100 %
11Beta-Hydroxy-Testosterone	5.00 %
Testosterone	2.00 %
4-Androstene-3,11,17-trione	0.83 %
DHT	0.50 %
11Beta-Hydroxy-Androstenedione	0.02 %
Androstenedione	<0.10 %
11-Oxo-Androsterone	0.02 %
11-Oxo-Etiocholanolone	<0.10 %
pregnenolone	0.02 %
Corticosterone	0.02 %
Cortisol	0.002 %
Estradiol	0.02 %

For research use only; not for use as a diagnostic.

