



Anti PACAP38 (1-15) (Human) Serum

Cat. No. YII-Y042-EX Lot No. 085171206

Description: This antiserum was raised in a rabbit by immunization with a porcine thyroglobulin (pTG) conjugate of synthetic PACAP (1-15) (human) peptide. The product vial contains 50 µL of the titled antiserum obtained by lyophilizing its 0.001 M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immuno-histochemistry or any other immunoreaction with the N-terminal portion of PACAP38 (human).

Immunogen: Synthetic PACAP38 (1-15) (human)-pTG conjugate **Host:** Rabbit

Amino Acid Sequence of PACAP38 (1-15) (human)¹⁾:

1 15

HSDGIFTDSY SRYRKQMAVK KYLAAVLGKR YKQRVKNK-NH₂

Product Form: Lyophilized unpurified serum **Size:** 50 µL

Reconstitution: Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN₃ 0.1%).

Storage: The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing-thawing should be avoided.

Suggested Working Dilution Range: 1:5,000-20,000 (final dilution ~1:140,000) for radioimmunoassay;

1: 500-3,000 for immunohistochemistry (frozen or paraffin sections). Optimal dilution should be determined by each laboratory for each application.

Specificity (based on radioimmunoassay): PACAP38 (1-15) (human) 100%, PACAP27 (human) 30%, PACAP38 (human) 6.7%, VIP (porcine) 0%, PHI (rat) 0%, PHI (human) 0%, secretin (human) 0%, helodermin 0%

Positive Control (immunohistochemistry): Rat colon

Species Tested: Rat²⁾, ginseng radix³⁾

REFERENCES:

- 1) A. Miyata, A. Arimura et al., Isolation of a novel 38 residue-hypothalamic polypeptide which stimulates adenylate cyclase in pituitary cells. *Biochemistry Biophysics Research Communication* 164:567-574, 1989
- 2) N. Yanaihara, T. Kanno et al., VIP- and PACAP-induced salivary chromogranin A secretion in the isolated perfused submandibular gland of rats. *Annals of the New York Academy of Sciences*, J. Fahrenkrug and S.I. Said (Ed), 921:218-225, 2000
- 3) N. Takashima, N. Yanaihara et al., Characterization of a PACAP-like immunoreactive component in red ginseng root. *Annals of the New York Academy of Sciences*, W.G. Forsmann (Ed) 865:561-565, 1998

FOR RESEARCH LABORATORY USE ONLY

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM

