



Anti Chromogranin A (359-389) (Rat) Serum

Cat. No. YII-Y290-EX **Lot No. 076371204**

Description: This antiserum was raised in a rabbit by immunization with a carrier free synthetic chromogranin A (359-389) (rat) peptide. The product vial contains 50 µL of the titled antiserum obtained by lyophilizing its 0.001 M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with chromogranin A related peptides.

Immunogen: Synthetic chromogranin A (359-389) (rat), carrier free **Host:** Rabbit

Amino Acid Sequence of Chromogranin A (359-389) (rat)¹⁾:

LE GEDDPDRSMK LSFRRARYGF RDPGPQLRR

Product Form: Lyophilized unpurified serum **Size:** 50 µL

Reconstitution: Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN₃ 0.1%).

Storage: The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing-thawing should be avoided.

Suggested Working Dilution Range: 1:2,000 (final dilution ~1:14,000) for radioimmunoassay^{2,3)}; 1: 1,000 for immunohistochemistry (frozen section)⁴⁾. Optimal dilution should be determined by each laboratory for each application.

Specificity (based on radioimmunoassay): Chromogranin A (359-389) (rat) 100%, pancreastatin (rat) 0%,

Positive Control (immunohistochemistry): Rat pituitary gland and adrenal gland^{5,6,7,8)}

Species Tested: Rat, dog, human

REFERENCES:

- 1) A. Iacangelo, H. Okayama, and L.E. Eiden, Primary structure of rat chromogranin A and distribution of its mRNA. FEBS Letter 227:115-121, 1988
- 2) S. Nagasawa, Y. Nishikawa et al., Distribution of chromogranin A-like peptides in the rat, Biomedical Research 16: 83-90, 1995
- 3) Y. Nishikawa, C. Yanaihara et al., Immunoreactive chromogranin A in rat tissue, plasma and urine, Peptide Chemistry 1994, Protein Research Foundation, Osaka, M. Ohno (Ed), pp389-392, 1995
- 4) T. Kanno, N. Yanaihara et al., Autonomic control of submandibular chromogranin A secretion in the anaesthetized rat. Biomedical Research 19: 411-414, 1998
- 5) Y. Nishikawa, T. Iwanaga et al., Distribution and existence pattern of chromogranin A in tissues studied by using region-specific antisera, Proceedings of the 11th Gut Hormone Conference, Japan Society of Gut Hormones (Ed), 1992, 2:p345-352
- 6) Y. Hashimoto, H. Ohki et al., Immunohistochemical demonstration of chromogranin A in endocrine organs of the rat and horse by use of region-specific antibodies, Japan Journal of Veterinary Research, 49: 3-17, 2001
- 7) N. Yanaihara, T. Kanno et al., VIP- and PACAP-induced salivary chromogranin A secretion in the isolated perfused submandibular gland of rats, Annals of the New York Academy of Sciences, 921:218-205, 2000
- 8) K. Ohshima, S. Nagasawa et al., Synthesis of rat chromogranin A fragments and their immunological properties, Peptide Chemistry 1991, Protein Research Foundation, Osaka, A. Suzuki (Ed), pp157-162, 1992

FOR RESEARCH LABORATORY USE ONLY

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM

