

# Product Information

## CF™ Dye Human Transferrin Conjugates

| Catalog # | Conjugate                  |
|-----------|----------------------------|
| 00081     | Human Transferrin, CF™488A |
| 00082     | Human Transferrin, CF™543  |
| 00083     | Human Transferrin, CF™568  |
| 00084     | Human Transferrin, CF™594  |
| 00085     | Human Transferrin, CF™640R |
| 00086     | Human Transferrin, CF™680R |
| 00087     | Human Transferrin, CF™750  |

**Unit Size:** 1 mg

### Storage and Handling

Store at -20°C upon arrival and protect from light. The lyophilized product is stable for at least six months from date of receipt when stored as recommended. Reconstitute in phosphate-buffered saline, add sodium azide to a final concentration of 2 mM, and store at 4°C. Briefly centrifuge the tube prior to use to remove any aggregates. This product contains human holo-transferrin and has tested negative for HIV and HBsAg, but should be treated as containing potentially infectious agents. Please handle and dispose of this product using universal laboratory safety precautions.


### Spectral Properties

| Conjugate                 | Abs <sub>max</sub> (nm) | Em <sub>max</sub> (nm) |
|---------------------------|-------------------------|------------------------|
| CF™488A Human Transferrin | 490                     | 515                    |
| CF™543 Human Transferrin  | 541                     | 560                    |
| CF™568 Human Transferrin  | 562                     | 583                    |
| CF™594 Human Transferrin  | 593                     | 614                    |
| CF™640R Human Transferrin | 642                     | 662                    |
| CF™680R Human Transferrin | 663                     | 682                    |
| CF™750 Human Transferrin  | 755                     | 777                    |

### Product Description

Transferrin is an iron-binding glycoprotein that delivers iron to cells by binding to the transferrin receptor when in its holo form (saturated with iron). The complex is subsequently internalized via clathrin-mediated endocytosis. Several factors, including endosomal pH and oxidoreductase activity, allow for the release of iron from transferrin. The apo-transferrin is then recycled back to the surface of the cell with the transferrin receptor via the endocytic recycling pathway and released to bind more iron.

Labeled human holo-transferrin is used for microscopic studies of the endosomal pathway and/or transferrin uptake. Biotium's holo-transferrin conjugates are labeled with a selection of our CF™ dyes, a series of next-generation fluorescent dyes developed at Biotium to have combined advantages in brightness, photostability, and water solubility compared to other fluorescent dyes.



**Figure 1. HeLa cells labeled with CF™488A Human Transferrin.** Cells were incubated at 4°C for 90 minutes with 25 ug/mL CF™488A Human Transferrin (green) and were either fixed immediately or placed at 37°C for 15 minutes prior to fixation. Cells were also stained with DAPI (blue). A) Surface labeling - cell surface view, B) surface labeling - cross-section of cells, C) trafficked transferrin - cross section of cells.

### Assay Protocol

The following is a protocol for labeling transferrin receptors on mammalian cells.

1. Reconstitute CF™ dye conjugated human transferrin in 1X PBS to a concentration of 1 mg/mL. For storage at 4°C, sodium azide can be added to a final concentration of 2 mM if it is compatible with your application.
2. Serum starve cells for one hour prior to labeling. Then, wash cells with cold PBS + 0.1% Bovine Serum Albumin (BSA). Add a final concentration of 25 ug/mL CF™ dye conjugated human transferrin in cold PBS + 0.1% BSA and incubate at 4°C for 90 minutes in the dark. Wash cells three times with cold PBS + 0.1% BSA.
3. A: For surface staining, immediately fix cells in 4% paraformaldehyde in 1X PBS for 15 minutes at room temperature (protected from light).  
B: For trafficking assays, add warm complete cell medium to the cells and incubate at 37°C for 5 minutes to 1 hour in the dark. Wash cells three times in cold PBS + 0.1% BSA prior to fixing the cells in 4% paraformaldehyde in 1X PBS for 15 minutes at room temperature (protected from light).
4. Wash cells twice with 1X PBS and process samples for imaging or subsequent immunostaining.

### Related Products

| Catalog # | Product Name  | Unit Size           |
|-----------|---|---------------------|
| 40061-T   | RedDot™2 Far Red Nuclear Counterstain, 200X in DMSO, Trial Size | 25 uL (15-20 tests) |
| 40046     | Hoechst 33342, 10 mg/mL in H <sub>2</sub> O                     | 10 mL               |
| 23001     | EverBrite™ Mounting Medium                                      | 10 mL               |
| 23002     | EverBrite™ Mounting Medium with DAPI                            | 10 mL               |
| 23003     | EverBrite™ Hardset Mounting Medium                              | 10 mL               |
| 23004     | EverBrite™ Hardset Mounting Medium with DAPI                    | 10 mL               |
| 23005     | CoverGrip™ Coverslip Sealant                                    | 15 mL               |
| 22005     | Mini Super <sup>HT</sup> Pap Pen 2.5 mm tip, ~400 uses          | 1 pen               |
| 22006     | Super <sup>HT</sup> Pap Pen 4 mm tip, ~800 uses                 | 1 pen               |

Please visit [www.biotium.com](http://www.biotium.com) to view our full selection of CF™ dye bioconjugates, including secondary antibodies, anti-tag and anti-hapten antibodies, phalloidin, alpha-bungarotoxin, lectins, Annexin V, and many other innovative products for life science research.

CF is a trademark of Biotium, Inc. CF dye technology is covered by pending US and international patents. Biotium products are high-quality reagents and materials intended for research purposes only. We welcome inquiries about licensing the use of our dyes, trademarks or technologies. Please submit inquiries by e-mail to [btinfo@biotium.com](mailto:btinfo@biotium.com).