

Product Information

Mix-n-Stain™ CF™660R antibody labeling kit, 1 labeling

Catalog Number: 92243, 50 - 100 ug labeling

Components and Storage:

Material	Quantity	Storage	Stability
CF660R dye (#92243A)	1 vial	-20 °C	Stable for 3 months from time of reception if stored as directed.
Mix-n-Stain reaction buffer, 10X (#99951)	1 vial of 30 uL		
Mix-n-Stain storage buffer (#99952)	1 vial of 300 uL		
Ultrafiltration vial (MWCO=10K) (#99956-1)	1 each		

Catalog Number: 92261, 20 - 50 ug labeling

Components and Storage:

Material	Quantity	Storage	Stability
CF660R dye (#92261A)	1 vial	-20 °C	Stable for 3 months from time of reception if stored as directed.
Mix-n-Stain reaction buffer, 10X (#99951-1)	1 vial of 15 uL		
Mix-n-Stain storage buffer (#99952-1)	1 vial of 150 uL		
Ultrafiltration vial (MWCO=10K) (#99956-1)	1 each		

Catalog Number: 92281, 5 - 20 ug labeling

Components and Storage:

Material	Quantity	Storage	Stability
CF660R dye (#92281A)	1 vial	-20 °C	Stable for 3 months from time of reception if stored as directed.
Mix-n-Stain reaction buffer, 10X (#99951-1)	1 vial of 15 uL		
Mix-n-Stain storage buffer (#99952-2)	1 vial of 60 uL		
Ultrafiltration vial (MWCO=10K) (#99956-1)	1 each		

Technical Summary

Abs/Em Maxima: 663/682 nm

Flow cytometry laser line: 633 or 635 nm

Microscopy laser line: 633 or 635 nm

Direct replacement for: Alexa Fluor® 660, Allophycocyanin (APC)

Alexa Fluor is a registered trademark of Invitrogen.
 CF™ dye technology is covered by pending US and international patents.

Product Application

This kit contains everything you need to rapidly label an antibody. Please select the labeling kit suitable for the amount of antibody you wish to label. The labeling procedure comprises simple mixing of your antibody and the optimally formulated dye in the reaction buffer provided, followed by a brief incubation. The resulting solution is ready for staining without further purification. The kit is suitable for labeling commercially available primary antibodies, either directly or after a simple antibody clean-up step. After labeling, the dye is covalently linked to the antibody.

CF™660R emits fluorescence at about 685 nm in the borderline spectral region between far-red and near-IR. Although its absorption maximum is around 660 nm, the dye can be sufficiently excited by the 633, 635 or 640 nm laser. When combined with other CF™ dyes of shorter wavelengths, CF™660R can serve as a useful long wavelength dye in multi-color detection. The dye is spectrally similar to Alexa Fluor® 660 but far superior to the latter in performance. The superior photostability and excellent brightness of CF™660R make the dye an ideal choice for confocal microscopy and other demanding applications.

Protocol

Before you begin:

a) Check the compatibility of your antibody with the antibody compatibility guide on the next page. Low levels of BSA, gelatin, Tris, glycerol or sodium azide in the antibody solution can be tolerated. However, if the amount of stabilizers are above the threshold concentrations, use either a protein A purification procedure or a commercial antibody clean-up kit, such as the Pierce antibody clean-up kit (cat#44600) to remove protein stabilizers. An antibody solution free of stabilizers produces better labeling results. If you need to remove just non-protein components such as Tris, glycine or glycerol, use the ultrafiltration vial provided in the kit to purify your antibody by following the steps below. If no clean-up is required, proceed to step b).

1. Add an appropriate amount of antibody to the ultrafiltration vial being careful not to touch the membrane. Spin the solution at max speed (14,000 rpm) in a microcentrifuge for a few minutes until the liquid is removed. Discard the liquid in the collection vessel.
2. Rinse the original antibody vial with an equal volume of 1X PBS and add to the ultrafiltration vial. Spin the vial at max speed until the liquid is removed. Resuspend the antibody in an appropriate amount of 1X PBS to derive a concentration of 0.5 - 1 mg/mL.

b) Use your antibody at a concentration of 0.5-1 mg/mL for optimal labeling. If the antibody is in a lyophilized form or is more concentrated, reconstitute or dilute the antibody in PBS. If your primary antibody is a commercial product with no quantity information, please contact the supplier. Resuspend the antibody in an appropriate amount of 1X PBS.

c) Warm up the Mix-n-Stain™ Reaction Buffer vial and the Mix-n-Stain™ Storage Buffer vial to room temperature before use. Centrifuge the vials briefly to concentrate the solutions to the bottom of the vials.

Procedure:

1. Dilute the 10X Mix-n-Stain™ Reaction Buffer to 1X in the antibody solution. Mix the solutions by pipetting up and down a few times.
2. Transfer the entire solution from Step 1 to the vial containing the CF™ dye (no need to measure the amount of the dye). Vortex the vial for a few seconds.
3. Incubate the vial in the dark for 30 minutes. The solution is now ready to use. The concentration of the CF dye-labeled antibody is approximately the amount of your starting antibody divided by the total volume (*i.e.*, ~100% labeling yield).
4. Dilute the labeled antibody solution with the provided Storage Buffer. Simply transfer the entire labeled antibody solution into the Storage Buffer and store at 4 °C. Recalculate the concentration of the antibody solution. Alternatively, you can aliquot the solution and store at -20 °C. Without repeated freeze-thaws, the shelf-life of the labeled antibody solution should be at least 6 months.




Figure 1. Absorption and emission spectra of CF™660C and CF™660R conjugated to goat anti-mouse IgG, respectively, in PBS.

Mix-n-Stain™ Compatibility with Common Antibody Storage Components

Component	Compatibility
BSA or gelatin	< 4 equivalent of antibody by weight
Glycerol	< 10%
Glycine	not compatible
Serum	not compatible
Sodium Azide	compatible
Tris	< 20 mM

* If your antibody contains storage buffer components above the compatibility threshold levels, purify your antibody.

** An antibody solution free of protein stabilizers produces the best results. Although low levels of BSA and gelatin can be tolerated, you may experience slightly higher background after staining if your antibody was labeled in the presence of these proteins.

*** If the antibody was labeled in the presence of BSA or gelatin, background staining can be greatly reduced by using blocking and wash solutions containing at least 1% BSA or gelatin, respectively.

Related Products

Cat.#	Product Name	Unit Size
23001	EverBrite™ Mounting Medium	10 mL
23002	EverBrite™ Mounting Medium with DAPI	10 mL
22010	10X Fish Gelatin Blocking Buffer	100 mL
22011	Fish Gelatin Powder	2 X 50 g
22010	Dry Milk Powder	4 X 25 g

Mix-n-Stain™ CF™ Dye Antibody Labeling Kits

Dye	Packaging	Cat. No.	Packaging	Cat. No.	Packaging	Cat. No.
CF™350	1 X (50-100 ug) labeling	92230	1 X (20-50 ug) labeling	92250	1 X (5-20 ug) labeling	92270
CF™405S	1 X (50-100 ug) labeling	92231	1 X (20-50 ug) labeling	92251	1 X (5-20 ug) labeling	92271
CF™405M	1 X (50-100 ug) labeling	92232	1 X (20-50 ug) labeling	92252	1 X (5-20 ug) labeling	92272
CF™488A	1 X (50-100 ug) labeling	92233	1 X (20-50 ug) labeling	92253	1 X (5-20 ug) labeling	92273
CF™555	1 X (50-100 ug) labeling	92234	1 X (20-50 ug) labeling	92254	1 X (5-20 ug) labeling	92274
CF™568	1 X (50-100 ug) labeling	92235	1 X (20-50 ug) labeling	92255	1 X (5-20 ug) labeling	92275
CF™594	1 X (50-100 ug) labeling	92236	1 X (20-50 ug) labeling	92256	1 X (5-20 ug) labeling	92276
CF™633	1 X (50-100 ug) labeling	92237	1 X (20-50 ug) labeling	92257	1 X (5-20 ug) labeling	92277
CF™640R	1 X (50-100 ug) labeling	92245	1 X (20-50 ug) labeling	92258	1 X (5-20 ug) labeling	92278
CF™647	1 X (50-100 ug) labeling	92238	1 X (20-50 ug) labeling	92259	1 X (5-20 ug) labeling	92279
CF™660C	1 X (50-100 ug) labeling	92239	1 X (20-50 ug) labeling	92260	1 X (5-20 ug) labeling	92280
CF™660R	1 X (50-100 ug) labeling	92243	1 X (20-50 ug) labeling	92261	1 X (5-20 ug) labeling	92281
CF™680	1 X (50-100 ug) labeling	92240	1 X (20-50 ug) labeling	92262	1 X (5-20 ug) labeling	92282
CF™680R	1 X (50-100 ug) labeling	92246	1 X (20-50 ug) labeling	92263	1 X (5-20 ug) labeling	92283
CF™750	1 X (50-100 ug) labeling	92241	1 X (20-50 ug) labeling	92264	1 X (5-20 ug) labeling	92284
CF™770	1 X (50-100 ug) labeling	92242	1 X (20-50 ug) labeling	92265	1 X (5-20 ug) labeling	92285

Biotium products are high-quality reagents and materials intended for research purposes only. Some products are potentially hazardous chemicals - please read the Material Safety Data Sheet for additional information regarding handling potentially hazardous chemicals. Several of Biotium products and product applications are covered by U.S. and international patents and pending patents. Our products are not available for resale or other commercial uses without a specific agreement from Biotium, Inc. We welcome inquiries about licensing the use of our dyes, trademarks or technologies. Please submit inquiries by e-mail to btinfo@biotium.com. All names containing the designation™ are trademarks of Biotium, Inc.

The manufacture, use, sale, offer for sale, or import of this product may be subject to one or more patents or pending applications owned or licensed by Biotium, Inc. The purchase price of this product includes a limited, non-transferable immunity and corresponding patent claims outside of the United States, to use only the purchased amount of the products and components of the product solely for the buyer's own internal research (whether the buyer is an academic or for-profit entity), in a manner consistent with the accompanying product literature. No right to use, sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for any other purpose or a Commercial Purpose, is hereby granted expressly, by implication or by estoppel. Commercial Purpose means any activity by a party for consideration and include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service including without limitation, reporting information or data resulting from use of the product or its components for a fee or other commercial consideration; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For products that are subject to multiple limited use label licenses, the most restrictive terms apply. This product is for research use only. Diagnostic uses require a separate license from Biotium, Inc. For information on purchasing a license to this product including for purposes other than research, contact Biotium, Inc., 3159 Corporate Place, Hayward, CA 94545, Tel: (510) 265-1027. Fax: (510) 265-1352.