

MONOCLONAL ANTIBODY

For research use only. Not for clinical diagnosis

Catalog No. CEC-044

Anti- SUMO2/3 antibody

BACKGROUND

SUMO is Ubiquitin-like protein that can be covalently attached to proteins as a monomer or a lysine-linked polymer. This post-translational modification on lysine residues of proteins plays a crucial role in a number of cellular processes such as nuclear transport, DNA replication and repair, mitosis and signal transduction.

Product type Primary antibody

Immunogen Recombinant GST-fused human SUMO3 (full length).

Source Culture supernatant

Purification Ion-exchange chromatography

Form Liquid

Presentation Purified monoclonal antibody in PBS, 50% Glycerol, 0.05%w/v ProClin300

 $\begin{tabular}{lll} \mbox{Concentration} & 1 \mbox{ mg/mL} \\ \mbox{Volume} & 100 \mbox{ μL} \\ \mbox{Label} & Unlabeled \\ \end{tabular}$

Specificity SUMO2 and SUMO3

Cross reactivity Human, Monkey, Mouse, Rat Other species have not been tested.

Storage Store below -20℃ (below -70℃ for prolonged storage)

Aliquot to avoid cycles of freeze/thaw.

Other -

Application notes Recommended use

WB, ICC, IHC Not tested for other applications.


Recommended dilutions

Western blotting, 1/1000 to 1/5000 Immunocytochemistry, 1/100 to 1/500 Immunohistochemistry, 1/100 to 1/500

Optimal dilutions/concentrations should be determined by the end user.

References 1) Saitoh et al.,(2006). Exp Cell Res., 312, 1418-1430.

* This antibody is used in ref.1.



Mouse primary neural progenitor cells

Fig.1 Immunocytochemistry/ Immunofluorescence – SUMO2/3 antibody (3H12), Mouse primary culture neuron




Fig.2 Western blot - SUMO1 antibody (4D12) and SUMO2/3 antibody (3H12), HeLa cell total extracts

For research use only, Not for diagnostic use.

