



POLYCLONAL ANTIBODY

For research use only, Not for diagnostic use.

Catalog No. ICA-TG2-RTP2

Anti Human ITIH4

[inter-alpha-trypsin inhibitor heavy chain family, member 4]

BACKGROUND

ITIH4 is the inter-alpha-trypsin inhibitor heavy chain family, member 4. The protein encoded by this gene is secreted into the blood, where it is cleaved by plasma kallikrein into two smaller forms. Expression of this gene has been detected only in liver.

Product type	Primary antibody
Immunogen	ICAntibodies™ encoding human ITIH4 full length nucleic acid sequence
Host Species	Sprague Dawley Rat
Fusion Partner	-
Clone Designation	-
Isotype	-
Source	Whole Serum
Purification	-
Formulation Buffer /Form	Liquid (Preservative : None)
Concentration	-
Volume	50 UL
Label	Unlabeled
Specificity	Human ITIH4
Cross species reactivity	Rat. Other species is not tested.
Storage Conditions	Store at -20°C (at -70°C for prolonged storage). Aliquot to avoid cycles of freeze/thaw. Storage Instructions : Prepare fresh working dilution before use. Avoid repeated freezing and thawing cycles (it is advisable to make aliquots before storage). Alternatively, add an equal volume of glycerol (ACS grade or better) for a final concentration of 50% and store at -20°C as a liquid. Note : Adding glycerol reduces the stated protein concentration and dilution range by one-half

Application notes	• Immunohistochemistry: 1/50
Recommended dilutions	• Flow cytometry: 1/100
	• ELISA: 1/500

Other applications have not been tested.
Optimal dilutions/concentrations should be determined by the end user.

References	-
-------------------	---

ANTIBODY CHARACTERIZATION




Fig.1 Immunohistochemistry: HeLa cells transfected with ICAfectin441 and plasmid encoding human ITIH4 were fixed, permeabilized and stained with rat anti-human ITIH4 Polyclonal Antibodies diluted 1/50 and Alexa488 goat anti rat IgG secondary antibody. Nuclei were counterstained with DAPI.




Fig.2 Flow cytometry: Untransfected (black) and transfected with ICAfectin 441 and plasmid encoding human ITIH4 (red) HeLa cells were fixed, permeabilized and stained with rat anti-human ITIH4 Polyclonal Antibodies diluted 1/100 and R-PE goat anti rat IgG secondary antibody.




Fig.3 ELISA: Lysates of untransfected and transfected HeLa cells with ICAfectin441 and plasmid encoding human ITIH4 were coated. Determination anti-human GPNMB Polyclonal IgG in the rabbit sera was performed by using HRP goat anti rabbit IgG secondary antibody.

RELATED PRODUCT:

Product Name	Antibody Specificity	Host Species	Application	Quantity	Maker	Cat#
Anti Human ITIH4 Polyclonal Antibody	Human / Rat	Brown Norway Rat	IHC / FCM / ELISA	50 UL	CAC	ICA-TG2-RTP1
Anti Human ITIH4 Polyclonal Antibody	Human / Rat	Sprague Dawley Rat	IHC / FCM / ELISA	50 UL	CAC	ICA-TG2-RTP2

Developed by In-Cell-Art 

For research use only, Not for diagnostic use.



COSMO BIO Co., LTD.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

URL: <http://www.cosmobio.co.jp>

e-mail: export@cosmobio.co.jp

[Outside Japan] Phone : +81-3-5632-9617

[国内連絡先] Phone : +81-3-5632-9610

FAX : +81-3-5632-9618

FAX : +81-3-5632-9619