



POLYCLONAL ANTIBODY

For research use only. Not for clinical diagnosis.

Catalog No. KZ-HS-P05

Anti Bovine Inhibin

BACKGROUND

Inhibin is an endocrine hormone produced in the ovaries, the testes and the placenta. This hormone has several functions in the body, with inhibin levels in women being linked to the menstrual cycle and playing a role in fetal development. Another hormone, activin, has an action in the body opposite to that of inhibin. Levels of these two hormones tend to fluctuate in both men and women in response to a number of cues which can include changes in hormone levels triggered by natural biological processes, environmental pressure, and other factors.


Product type	Primary antibodies
Host	Goat
Form	Serum
Volume	50 µl
Antigen	30 kDa bovine inhibin
Cross reactivity	Bovine, human, ovine, rat, mouse
Application notes	Recommended dilutions

RIA : 1/10,000 - 1/500,000

IHC : 1/1,000 - 1/5,000

WB : 1/1,000 - 1/5,000

Neutrization : n/d



Dose-response curves of various inhibin-related proteins in the time-resolved fluoroimmunoassays for inhibin A. Materials tested were 32-kDa inhibin A (●), activin A (▲), activin AB (■), activin B (○), pro-αC (▼) purified from bovine follicular fluid, and human recombinant inhibin B (□). Modified from Kaneko *et al.* (2002a).

Storage

Store below -20°C (below -70°C for prolonged storage).
Aliquot to avoid cycles of freeze/thaw.

References

- 1) Hasegawa Y, Madarame H & Ibuki Y (1995) Inhibin and activin: Novel regulators for gonadal function and sexual maturation. Eds. Hibi I and Tanaka T, in: *Sexual Differentiation and Maturation*, P. 139-154, Ares Serono Symposia Publications
- 2) Kaneko H, Taya K, Watanabe G, Noguchi J, Kikuchi K, Shimada A & Hasegawa Y (1997) Inhibin is involved in the suppression of FSH secretion in the growth phase of the dominant follicle during the early luteal phase in cows. *Domestic Ani Endocrinol.* **14**: 263-271. PMID: [9260064](#).
- 3) Hasegawa Y, Madarame H, Yoshida S, Kaneko H, Abe Y, Mizunuma H & Ibuki Y (1997) Two-site immunoassay for native inhibin A. Eds. Aono T, Sugino H and Vale W. in: *Inhibin, Activin and Follistatin*, P. 104-117, Springer-Verlag
- 4) Kaneko H, Noguchi J, Kikuchi K, Akagi S, Shimada A, Taya K, Watanabe G & Hasegawa Y (2001) Production and endocrine role of inhibin during the early development of bull calves. *Biol Reprod.* **65**: 209-215. PMID: [11420242](#)
- 5) Kaneko H, Noguchi J, Kikuchi K, Todoroki J & Hasegawa Y (2002) Alterations in peripheral concentrations of inhibin A in cattle studied using a time-resolved immunofluorometric assay: Relationship with estradiol and follicle-stimulating hormone in various reproductive conditions. *Biol Reprod.* **67**: 38-45. PMID: [12079997](#)
- 6) Kaneko H and Hasegawa Y (2007) Application of time-resolved fluorometry to immunoassays for bovine reproductive hormones. *Ani Sci J.* **78**: 7-15.

For research use only. Not for clinical diagnosis.



COSMO BIO Co., LTD.

【JAPAN】

TOYO EKIMAE BLDG. 2-20, TOYO 2-CHOME,
KOTO-KU. TOKYO 135-0016, JAPAN
Phone: +81-3-5632-9610
FAX: +81-3-5632-9619
URL: <https://www.cosmobio.co.jp/>



COSMO BIO USA

【Outside Japan】

2792 Loker Ave West, Suite 101
Carlsbad, CA 92010, USA
email: info@cosmobioussa.com
Phone/FAX: (+1) 760-431-4600
URL: www.cosmobioussa.com