Catalog No. TIP-SN-P09

Anti α-Synuclein (131-140)

BACKGROUND

Alpha-Synuclein, a 140-amino acid protein abundantly expressed in presynaptic terminals, is known as a component of intraneuronal or glial inclusions observed in cases of Parkinson's disease (PD), Dementia with Lewy bodies (DLB) and Multiple system atrophy (MSA). Alpha-Synuclein is a natively unfolded protein, however, fibrillization or conformational change of Alpha-synuclein is central in the pathogenesis of Alpha-synucleinopathies. The amino-terminal region of Alpha-synuclein consists of seven imperfect repeats, each 11 amino acids in length, with the consensus sequence KTKEGV. The repeats partially overlap with a hydrophobic region (amino acids 61-95). The carboxy-terminal region (amino acids 96-140) is negatively charged. These antibodies are powerful tools for biochemical and IHC analyses of neurodegenerative diseases and for evaluation of conformational changes of Alpha-synuclein.

Product type Primary antibody

ImmunogenCEGYQDYEPEA (α-syn131-140)Raised inRabbit (New Zealand White)

Source Anti-serum

Purification ammonium sulfate precipitation

Form Liquid containing 50% glycerol/PBS with 0.05% NaN3 as a preservative.

 Concentration

 Volume
 50 uL

 Label
 Unlabeled

 Specificity
 α-Syn

Cross reactivity Human, Mouse

Storage Store below -20°C. (below -70°C for prolonged storage).

Aliquot to avoid cycles of freeze/thaw.

Application notes Recommended

dilutions

ELISA: 1/1000-1/2000Western Blot: 1/1000

•Immunohistochemistry:1/1000-1/2000

Other applications have not been tested. Optimal dilutions/ concentrations

should be determined by the end user.

References 1)Masami Masuda, et al. Inhibition of a-synuclein fibril assembly by small molecules:

Analysis using epitope-specific antibodies. FEBS Letters (2009) 583, 787-791. PMID

19183551

2) Motokuni Yonetani, et al. Conversion of wild-type alpha-synuclein into mutant-type fibrils and its propagation in the presence of A30P mutant. Journal of Biological

Chemistry (2009) 284, 7940-7950.

PMID 19164293

For research use only, Not for diagnostic use.



COSMO BIO CO., LTD.

(JAPAN)

TOYO EKIMAE BLDG. 2-20, TOYO 2-CHOME, KOTO-KU. TOKYO 135-0016, JAPAN Phone: +81-3-5632-9610

FAX: +81-3-5632-9619 URL: https://www.cosmobio.co.jp/



COSMO BIO USA

[Outside Japan] 2792 Loker Ave West, Suite 101 Carlsbad, CA 92010, USA email: info@cosmobiousa.com Phone/FAX: (+1) 760-431-4600 URL: www.cosmobiousa.com

RELATED PRODUCTS:

Product Name	Quantity	Maker	Cat#
Anti α-Synuclein (1-10)	50 uL	CAC	TIP-SN-P01
Anti α-Synuclein (11-20)	50 uL	CAC	TIP-SN-P02
Anti α-Synuclein (21-30)	50 uL	CAC	TIP-SN-P03
Anti α-Synuclein (31-40)	50 uL	CAC	TIP-SN-P04
Anti α-Synuclein (41-50)	50 uL	CAC	TIP-SN-P05
Anti α-Synuclein (51-60)	50 uL	CAC	TIP-SN-P06
Anti α-Synuclein (61-70)	50 uL	CAC	TIP-SN-P07
Anti α-Synuclein (75-91)	50 uL	CAC	TIP-SN-P08
Anti α-Synuclein (9 Antibodies Set)	9*10 uL	CAC	TIP-SN-SET