

Cosmo Bio Co., Ltd

TOYO EKIMAE BLDG, 2-20,TOYO 2CHOME, KOTO-KU,TOKYO 135-0016,JAPAN

TEL: +81-3-5632-9617 FAX: +81-3-5632-9618

URL: http://www.cosmobio.co.jp/e-mail: export@cosmobio.co.jp

国内連絡先

TEL: 03-5632-9610 FAX: 03-5632-9619

Catalog No.CBX00495

Mouse monoclonal antibody Anti-Human TAF9L

■ Formulation

Mouse monoclonal anti-human **TAF9L** antibody in PBS (3.0 mM KCl, 1.5 mM KH₂PO₄, 140 mM NaCl, 8.0 mM Na₂HPO₄ (pH 7.4)) containing 1% bovine serum albumin (BSA) and 0.05% sodium azide (NaN₃).

Antibody concentration

 $100 \mu g/ml (1.0 ml)$

■Storage

Store at 2-8°C for up to one year. We recommend storing at -20°C for long-term storage. Avoid repeat freezing and thawing cycles.

Preparation

This antibody was purified using protein G column chromatography from culture supernatant of hybridoma cultured in a medium containing bovine IgG-depleted (approximately 95%) fetal bovine serum.

Sterility

Filtered through a 0.22 µm membrane.


Applications

WB, IC, Dot Blot

Disposal

This antibody solution contains sodium azide (NaN₃) as a preservative. There is a potential hazard that NaN₃ reacts with copper or lead to produce an explosive compound. For safe disposal, the vial has to be washed thoroughly with water.

Lot No. 3365C4a-1 Clone No. 3365C4a Antibody class: IgG2b Immunogen: Recombinant



■ Safety warnings and precautions

Caution must be taken to avoid contact with skin or eyes. In such a case, rinse thoroughly at once with water. Do not ingest, inhale, or swallow. Seek medical attention immediately.

Wear appropriate protective clothing such as laboratory overalls, safety glasses and gloves.

It is strongly advised that this product should be handled by people who have been well trained in laboratory techniques and that it is handled with care pursuant to the principles of good laboratory practice. All chemicals are deemed potentially harmful.

The vial is prone to fall over. Use caution, especially when the lid is off.



Catalog No. CBX00495

Mouse monoclonal antibody Anti-Human TAF9L


Background

Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a protein that is similar to one of the small subunits of TFIID, TBP-associated factor 9, and is also a subunit of TFIID. TAF9 and TAF9b share some functions but also have distinct roles in the transcriptional regulatory process. [NCBI Entrez Gene Summary]

Recommended condition


WB: 0.2-2 μg/ml IC: 2-100 μg/ml

Application



Detection of TAF9L by Western blot. Samples: Whole cell lysate from human HEK293 (H, 25 μ g) , mouse NIH3T3 (M, 25 μ g) and rat F2408 (R, 25 μ g) cells. [Lot No. 3365C4a-1]

Predicted molecular weight: 27 kDa



Immunostaining analysis in HeLa cells. HeLa cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100 in PBS. The cells were immunostained with anti-TAF9L mAb. [Lot No. 3365C4a-1]