



## GA-BSA/ Glycolaldehyde-BSA

<b>Product Description</b>	Glycolaldehyde (33 mM) was incubated with bovine serum albumin (BSA) (2 mg/ml) at 37°C for 7 days in PBS (pH 7.4), and dialyzed against PBS.
<b>Volume</b>	200 ul
<b>Concentration</b>	1 mg/ml
<b>Storage</b>	Store below -20°C (below -70°C for prolonged storage). Aliquot to avoid cycles of freeze/thaw.

- References**
1. Nagai R., Hayashi CM., Xia L., Takeya M., Horiuchi S: Identification in human atherosclerotic lesions of GA-pyridine, a novel structure derived from glycolaldehyde-modified proteins. J Biol Chem. 277, 48905-48912 (2002) PMID: [12377783](#)
  2. Nagai R., Matsumoto K., Ling X., Suzuki H., Araki T., Horiuchi S: Glycolaldehyde, a reactive intermediate for advanced glycation endproducts, plays an important role in the generation of an active ligand for the macrophage scavenger receptor. Diabetes 49, 1714-1723, (2000) PMID: [11016456](#)

### Characterization




Fig.1 Immunoreactivity of CEL(CEL-SP) monoclonal antibody to CEL-BSA and CML-BSA

## ELISA protocol

### Coating

- 1) Distribute 100 µl / well of the sample solution (1 µg/mL in PBS) to 96 well microtiter plates (Thermo, MaxiSorp).
- 2) Incubate the plates 2h at RT or overnight at 4 degrees.
- 3) Discard the supernatant of sample solution.
- 4) Wash the plates three times with washing.buf.(PBS/0.05%Tween 20)

### Blocking

- 1) Distribute 200 µl / well of 0.5% gelatin-PBS to 96 well microtiter plates
- 2) Incubate the plates 1h at RT.
- 3) Discard the the supernatant of 0.5% gelatin-PBS
- 4) Wash the plates three times with washing.buf.(PBS/0.05%Tween 20)

### Primary antibody

- 1) Distribute 100 µl / well of Primary antibodies diluted with washing buf. as suggested in the APPLICATIONS to each well.
- 2) Incubate the plates 1h at RT.
- 3) Discard the supernatant of Primary antibody solution.
- 4) Wash the plates three times with washing.buf.(PBS/0.05%Tween 20)

### Secondary antibody

- 1) Distribute 100 µl / well of secondary antibodies (HRP-anti mouse IgG) diluted with washing buf. as suggested in the APPLICATIONS to each well.
- 2) Incubate the plates 1h at RT.
- 3) Discard the supernatant of secondary antibody.
- 4) Wash the plates three times with washing.buf.(PBS/0.05%Tween 20)

### OPD color reaction

- 1) Reaction for 2-10 minutes at RT..
- 2) Distribute 100 µL / well of 2M H<sub>2</sub>SO<sub>4</sub> to each well and stop enzyme reaction.
- 3) After gentle mixing, determine the absorbance at 492 nm of each well by a spectrophotometer.

## RELATED PRODUCTS:

Product Name	Quantity	Maker	Cat#
Anti N <sup>ε</sup> -(carboxymethyl) lysine [CML] (2G11) Monoclonal Antibody	100 ul	CAC	AGE-M01
Anti N <sup>ε</sup> -(carboxyethyl) lysine [CEL] (CEL-SP) Monoclonal Antibody	100 ul	CAC	AGE-M02
Anti GA-pyridine (2A2) Monoclonal Antibody	100 ul	CAC	AGE-M03
Anti N <sup>ω</sup> -(carboxymethyl) arginine [CMA] (3F5) Monoclonal Antibody	100 ul	CAC	AGE-M04
CML-BSA/Nε-(carboxymethyl) lysine-BSA	200 ul	CSR	AGE-GP01
CEL-BSA/Nε-(carboxyethyl) lysine-BSA	200 ul	CSR	AGE-GP02
GA-BSA/Glycolaldehyde-BSA	200 ul	CSR	AGE-GP03
Ribose-gelatin	500 ul	CSR	AGE-GP04
Mild-AGE-BSA	200 ul	CSR	AGE-GP05

*For research use only, Not for diagnostic use.*



**COSMO BIO Co., LTD.**

【JAPAN】

TOYO EKIMAE BLDG. 2-20, TOYO 2-CHOME,  
KOTO-KU. TOKYO 135-0016, JAPAN  
Phone: +81-3-5632-9610  
FAX: +81-3-5632-9619  
URL: <https://www.cosmobio.co.jp/>



**COSMO BIO USA**

【Outside Japan】

2792 Loker Ave West, Suite 101  
Carlsbad, CA 92010, USA  
email: [info@cosmobiousa.com](mailto:info@cosmobiousa.com)  
Phone/FAX: (+1) 760-431-4600  
URL: [www.cosmobiousa.com](http://www.cosmobiousa.com)