



biosciences incorporated P.O. Box 58537

Salt Lake City, UT 84158-0537

Tel: 801-588-0455 Fax: 801-588-0497

echelon@echelon-inc.com

For Investigational Use Only

## **Shuttle PIPTM Carrier**

Intracellular delivery of phosphoinositides

**Product Number: P-9C2** 

**Contents:** 

Catalog #<br/>P-9C2Description<br/>Histone H1Molecular Weight<br/>~26,230Quantity<br/>2 X 50 nmoles

**Storage:** Carrier 2 is lyophilized. Protect from moisture and store at -20 °C until reconstituted. Reconstitute with water or other aqueous solutions and store at 4°C after reconstituting for up to 3 months. Multiple freeze thawing is not recommended. *Note: phosphate buffers are not recommended and may alter complex formation with phosphoinositides.* We do not recommend storing carriers and PIPs together as complexes.

**Use**: Carriers are used to deliver phosphoinositide polyphosphates into living cells. This carrier has successfully delivered the following phosphoinositides into cells:  $PtdIns(4,5)P_2$ ,  $PtdIns(3,4)P_2$ , and  $PtdIns(3,4,5)P_3$ , and their fluorescent derivatives.

## **References:**

- 1. Ozaki, S., DeWald, D.B., Shope, J.C., Chen, J., Prestwich, G.D. Intracellular delivery of phosphoinositides and inositol phosphates using polyamine carriers. *Proc Natl Acad Sci U S A* **97**, 11286-91 (2000).
- 2. Wang, Y.J., Wang, J., Sun, H.Q., Martinez, M., Sun, Y.X., Macia, E., Kirchhausen, T., Albanesi, J.P., Roth, M.G., Yin, H.L. Phosphatidylinositol 4 phosphate regulates targeting of clathrin adaptor AP-1 complexes to the Golgi. *Cell* **114**, 299-310 (2003).
- 3. Maffucci, T., Brancaccio, A., Piccolo, E., Stein, R.C., Falasca, M. Insulin induces phosphatidylinositol-3-phosphate formation through TC10 activation. *Embo J* 22, 4178-89 (2003).
- 4. Larsen, M., Hoffman, M.P., Sakai, T., Neibaur, J.C., Mitchell, J.M., Yamada, K.M. Role of PI 3-kinase and PIP3 in submandibular gland branching morphogenesis. *Dev Biol* **255**, 178-91 (2003).
- 5. Weiner, O.D., Neilsen, P.O., Prestwich, G.D., Kirschner, M.W., Cantley, L.C., Bourne, H.R. A PtdInsP(3)- and Rho GTPase-mediated positive feedback loop regulates neutrophil polarity. *Nat Cell Biol* 4, 509-13 (2002).

Echelon Biosciences products are sold for research and development purposes only and are not for diagnostic use or to be incorporated into products for resale without written permission from Echelon Biosciences. Materials in this publication, as well as applications and methods and use, may be covered by one or more U.S. or foreign patents or patents pending. We welcome inquiries about licensing the use of our trademarks and technologies at busdev@echelon-inc.com.

TDS P-9C2 Rev: 2 (12/02/03)