

# PRODUCT INFORMATION **Biotin Labeled Lectin**

Catalog Number: BA-3101-2

Pure Wisteria floribunda lectin (WFA) from Japanese wisteria, Biotin Description:

conjugated.

Lot Number:

Protein

2 mg purified WFA Biotin/vial. Reconstitute with Buffer to a final

Concentration: concentration of 1mg/ml if lyophilized.

(Based on OD 280)

Carbohydrate Specificity:

N-Acetylgalactosamine

Inhibitory Carbohydrate: N-Acetylgalactosamine >> Lactose >> Galactose

Activity: 15-30 μg/ml will agglutinate type A<sub>1</sub> human erythrocytes. Less than

0.5 µg/ml will agglutinate neuraminidase treated erythrocytes.

Buffer: 0.01M Phosphate - 0.15M NaCl, pH 7.2 - 7.4.

**Chemical Used for** Conjugation:

Biotinyl N - hydroxysuccinimide ester (BNOHSE).

Storage: Store lyophilized powder refrigerated at 5-8°C or frozen. Store liquid

frozen in aliquots. Avoid freeze-thaw cycles.

Stability: The lyophilized material is stable for several years when stored frozen.

After reconstitution the material is stable for at least 1 year when stored frozen in aliquots with 0.05% sodium azide added as a preservative.

Caution: Refer to the enclosed MSDS for information regarding lectins. The

aluminum seals have sharp edges and the vial itself may have cracks

which can cause lacerations. Use caution when opening the vial.

See reverse side. Procedure for Use:

References: Torres, B.V., et al. (1988) Arch. Biochem. Biophys. 262:1-11.

Tsuda, M., et al. (1975) Gann: Japanese J. Cancer Res. 66:513-521.

Tel: 650-342-3296

Wong, K.H., et al. (1986) K. Clin. Micro. 23:407-410.

**LY** LABORATORIES, INC.

107 North Amphlett Blvd. San Mateo, CA 94401 650-342-2648 Fax: 1-800-821-0044 Orders: (Outside CA only)

# **General Procedure**

The following is a general Procedure and Trouble-Shooting Guide. The information is provided only for your convenience. The success of your experiments are not guaranteed by EY Laboratories. Inc.

- Wash and block tissue section or blot. EY Laboratories, Inc. recommends that 1% purified Bovine Serum Albumin (BSA) or defatted milk powder be used for blocking to prevent nonspecific binding. Do not use serum products, they contain glycoproteins which may lead to high levels of non specific background. After blocking, rinse briefly with recommended Buffer.
- Dilute Biotin Labeled Lectin to a concentration of 5-50 µg/ml using recommended Buffer. Incubate section or blot for 30-90 minutes at room temperature in a moist chamber. Slightly longer incubation times may be required if incubation is done at 2-8°C. Rinse 3 times, 5 minutes each time with recommended Buffer.
- Dilute and incubate Avidin Conjugate according to manufacturer directions.

**Notes:** Inhibition of lectin binding may be accomplished by using one of two procedures:


- A. Before proceeding to Step #3 incubate lectin treated section or blot with inhibitory carbohydrate for 30-60 minutes at room temperature. NOTE: Complete inhibition may
- Preincubate diluted Biotin Labeled Lectin with inhibitory carbohydrate for 30-60 minutes at room temperature before applying to section or blot.

#### TROUBLE SHOOTING GUIDE

Problem	Cause	Solution
Weak or no Staining	Low concentration of specific	Causes #1 - #4
	oligosaccharide on sample.	Increase incubation time.
	2. Low concentration of lectin conjugate.	b. Increase concentration of sample (on
	3. Low concentration of avidin conjugate.	blot) lectin conjugate and/or avidin
	<ul><li>4. Insufficient incubation time.</li><li>5. Inappropriate treatment of sample prior</li></ul>	conjugate. a. Treat section or blot with a different
	to labeling.	blocking reagent.
High Background	Lectin conjugate and/or avidin conjugate	Decrease concentration of respective
	is too concentrated.	reagents.
		b. Shorten incubation times.
	<ol><li>Insufficient washing.</li></ol>	<ul> <li>a. Perform multiple washings and</li> </ul>
		prolong washing time.
	<ol><li>Insufficient blocking.</li></ol>	<ul> <li>a. Treat section or blot with a different</li> </ul>
		blocking reagent.
	4. Sample contains endogenous enzymatic	a. Determine if sample contains activity
	activity.	which would give background staining
T I		in the absence of the avidin conjugate.  a. Perform control reactions.
Unexpected	Market 1	
Staining Pattern	Multiple causes	b. Use other cytochemical technique
rattern		to prove or disprove the findings.

Y LABORATORIES, INC.

650-342-3296 Tel: 650-342-2648 107 North Amphlett Blvd. Fax: 1-800-821-0044 Orders: San Mateo, CA 94401 (Outside CA only)



# **MATERIAL SAFETY DATA SHEET**

Effective Date: March 31, 2006 Revision 5 Page 1 of 2

## PRODUCT IDENTIFICATION

Name: Purified proteins and enzymes labeled with D-Biotin.

Catalog BAP-01. BA-1102 to BA-9000. BAF-001 to BAF-2354. BAL-1104 to BAL-4701. Number(s): BA-01 to BA-013, BA-108, BA-109, BA-111, BA-118, BA-119, BA-120, BA-121,

BAT-2100 to BAT-2701.

Formula: Complex polypeptides labeled with D-Biotin

Synonyms: Protein A. Lectins, Secondary and Monoclonal Antibodies, Horseradish

Peroxidase, Alkaline Phosphatase, Lactoperoxidase, Ferritin, and Urease labeled

NOTE: D-Biotin is also known as vitamin H.

#### **EMERGENCY INFORMATION**

EY Laboratories, Inc. 107 North Amphlett Blvd. San Mateo, CA 94401

# **EMERGENCY PHONE:** 650-342-3296

#### HAZARDOUS COMPONENTS

Specific protein (s) as listed on the vial label. Solutions are at a concentration generally greater than 0.5mg protein/ ml. Powders are >>95% pure protein unless otherwise indicated on the vial label. Biological activity of these labeled proteins will vary. Vitamin H is an essential vitamin, required in very low amounts. The concentration of bound biotin is less than 10% of the protein amount (w/w). All solutions contain less than 0.05% sodium azide as a preservative.

## **HEALTH HAZARD INFORMATION**

EXPOSURE LIMITS: None established. The toxicological properties of these products have not

been thoroughly investigated. Care should be taken when handling any of

**FFFFCTS OF** May cause localized eye, skin, or mucous membrane irritation. Some

OVEREXPOSURE: sensitive individuals may develop a chronic allergic reaction with exposure.

The known effects are due to the protein.

**ROUTES OF** Inhalation of powders and skin contact with liquids are the primary routes of **EXPOSURE:** 

exposure. Care should be taken to avoid the formation of aerosols when

handling any of the solutions.

#### PHYSICAL CHARACTERISTICS

APPEARANCE: Powders are white to brown. Solutions will be clear to dark brown or red. SOLUBILITY: Powders are completely soluble in many biological buffers and water.

All liquids are completely miscible in water and biological buffers.

LABORATORIES, INC.

197 North Amphlett Blvd. San Mateo, CA 94401

Tel: 650-342-3296 Fax: 650-342-2648 1-800-821-0044 Orders: (Outside CA only)

## MSDS for Biotin labeled Purified Proteins Continued - page 2 of 2.

FIRE AND EXPLOSION HAZARDS Not considered to be a fire hazard.

**EXTINGUISHING MEDIA:** Water spray or CO2.

SPECIAL FIRE FIGHTING NOTE: None required. NOTE:

Most solutions contain less than 0.05% sodium azide as a preservative. Azide may react with lead and copper plumbing to form explosive metal azides. Flush with copious amounts of water when disposing material in the

sink.

REACTIVITY DATA

STABILITY: Stable. Decomposition products are not known to be

hazardous.

HAZARDOUS POLYMERIZATION: Will NOT occur.

INCOMPATIBILITY: None known. (Lead and copper may react with sodium

SPILL / LEAK PROCEDURES

MATERIAL RELEASE / Avoid contact with powder or liquid. Clean up spill with a paper towel SPILL:

soaked in household bleach. Do not allow solutions to dry on environmental surfaces. Wash affected area with detergent after the area

has been treated with bleach.

Incinerate, autoclave, or dispose of paper waste in accordance with all WASTE DISPOSAL:

Local, State, and Federal regulations. Due to the small quantities of material involved these products are generally not considered to be environmental hazards. All of these proteins are fully biodegradable.

## **EMERGENCY FIRST AID PROCEDURES**

May be harmful if swallowed, inhaled, or allowed to absorb through the skin. Wash contacted area with water for 15 minutes. If inhaled remove to fresh air. Report exposure to the appropriate safety official. Consult a physician if irritation occurs or if there is any indication of an allergic response, such as watering eyes, sneezing, or difficulty breathing.

### SPECIAL HANDLING PRECAUTIONS

VENTILATION: No special ventilation is required but it is recommended to handle these

reagents in a fume hood when possible.

EYE PROTECTION: Not required under most circumstances but recommended as a safety

RESPIRATORY Recommended as a safety precaution, specifically when working with

powders. An approved respirator may be required for those individuals PROTECTION:

already known to be sensitive to these materials.

PROTECTIVE GLOVES: Required when handling any of these materials.

#### SPECIAL PRECAUTIONS

This material is for research and experimental application only. It is not intended for food, drug, household, agricultural, or cosmetic use. All materials should be handled only by technically qualified individuals experienced with working with potentially hazardous chemicals. The above information is correct to the best of our knowledge. The user should make independent decisions regarding completeness of the information, based on all sources available. EY Laboratories, Inc. shall not be held liable for any damage resulting from handling or contact with the above product.

Y LABORATORIES, INC.

107 North Amphlett Blvd. San Mateo, CA 94401

Tel: 650-342-3296 Fax: 650-342-2648 1-800-821-0044 Orders: (Outside CA only)