

# Silica Hydrogel Kit™

## User Guide

HR2-310

Gels are a very efficient media for growing macromolecular crystals. Silica gels in particular have the advantage in that they are stable, usable over a wide range of temperature (0–60°C), and are compatible with a wide variety of precipitants and additives used for crystal growth. The Hampton Research Silica Hydrogel™ kit provides you with the material required to make gels at room temperature which polymerize rapidly and form a porous matrix with a pH of 7.0. The neutral pH allows one to use the gel with minimal influence on the pH of the precipitant.

### Suggested Uses

- To reduce excessive nucleation
- To reduce excessive precipitate
- To grow fewer, larger crystals
- To change the morphology of the crystal
- To slow the crystal nucleation and/or growth of crystals and to alter the kinetics of the experiment.

These effects and their extent will vary with each macromolecule and precipitant/crystallization system. It should be noted that in some instances the use of a gel for crystal growth may have a detrimental effect under the selected crystallization conditions. One should seek to optimize conditions for gel growth (temperature, precipitant type/concentration, etc.).

### Sample Preparation


The sample should be as pure as is practically possible (>95%) and free of amorphous and particulate material. Remove amorphous material by centrifugation or micro-filtration prior to use.

Recommended sample concentration is 5 to 25 mg/ml. The sample should be free of any unnecessary additives. Ideally, the initial screen should be performed with a sample which has been dialyzed against water. Ligands, ions, reducing agents, or other additives may be present as required by the sample for solubility, stability, or activity.

### Using the Silica Hydrogel - Sitting Drop Methodology

1. The gel is created by mixing equal parts of the sodium silicate solution (Green Top Tube) with dilute acetic acid (Blue Top Tube). This can be accomplished by placing the entire 0.5 ml of the 2.8% (v/v) sodium silicate solution (Green Top Tube) into the dilute acetic acid solution (Blue Top Tube), or mix any custom volume you might require in a separate container as long as the ratio of sodium silicate to acetic acids is 1:1 (i.e. 200 µl plus 200 µl). **Do not add the acetic acid into the sodium silicate!** It is very important that the sodium silicate be added to the acetic acid solution. Figure 1.


**Figure 1**  
Mix the Sodium Silicate with the Acetic Acid.



2. Mix by vortexing for 30 seconds. This is your gel stock. It will polymerize in 5 to 10 minutes. The gel will have a neutral pH of 7.0.

3. Pipet 1 - 5 µl of the gel onto a Micro-Bridge or the post of the Cryschem plate. Allow to polymerize. This will require approximately 5 to 10 minutes. Allow one hour for a complete cure. A small amount of water may appear on the surface of the gel after cure. Remove this water with a Paper Wick or small piece of filter paper. Seal the reservoir with a cover slide or Crystal Clear Sealing Tape while you are waiting for the gel to polymerize. The gel will dehydrate if left open.

**Figure 2**  
Cross section of a reservoir in the Cryschem plate.



4. Add 1 - 5 µl of the precipitant you intend to place in the reservoir to the TOP (not the side) of the gel drop. Figure 2. Allow this precipitant to be completely absorbed into the gel. Complete absorption will require one hour for dilute precipitants and as long as eight hours for high concentrations of PEG and MPD. Again, keep the reservoir covered during this time.

5. Place 2 - 10 µl of your protein/peptide/small molecule solution onto the gel. Allow to soak in completely. This will require approximately one hour. Keep the reservoir covered during this time.

6. Place 0.7 to 1.0 ml of precipitant into the reservoir and seal the plate with Crystal Clear Sealing Tape or a cover slide.


### Using the Silica Hydrogel - Liquid / Gel Diffusion

See Sitting Drop Vapor Diffusion for inscriptions on preparing the gel solution.

1. Place 2 - 20 µl of gel into a small culture tube. Allow the gel solution to polymerize and cure for approximately one hour.


2. Add 2 - 20 µl of protein to the polymerized gel. Allow the material to be completely absorbed. This should require approximately one hour.

**Figure 4**  
Typical observations in a crystallization experiment



3. Add 4 - 40  $\mu$ l of precipitant to the top of the gel after the protein has been completely absorbed. Seal with parafilm or wax. See Figure 3 below.

**Figure 3**  
Liquid / Gel Diffusion



### Using the Silica Hydrogel - Liquid / Gel / Liquid Diffusion

See Sitting Drop Vapor Diffusion for instructions on preparing the gel solution.


1. Using a capillary or caraway tube (Fisher Scientific part number 02-668-35) place a small amount of gel into the center of the tube. Use sufficient gel to create a plug. The length of the plug should be approximately 2 mm. Longer plugs will slow diffusion between precipitant and sample but will also dilute your sample as it diffuses. Allow the gel to polymerize.

2. Place 1 - 20  $\mu$ l of your sample on one side of the gel plug using a thin tipped sequencing gel loading pipet tip.

3. Place 1 - 20  $\mu$ l of your precipitant on the opposite side of the gel.

4. Seal the ends of the tube with parafilm, mounting clay, or wax.

**Figure 4**  
Liquid / Gel / Liquid Diffusion



### Examine the Drop

Carefully examine the drops under a stereo microscope (10 to 100x magnification) immediately after setting up the screen. Record all observations and be particularly careful to scan the focal plane for small crystals. Observe the drops once each day for the first week, then once a week there after. Records should indicate whether the drop is clear, contains precipitate, and or crystals. It is helpful to

describe the drop contents using descriptive terms. Adding magnitude is also helpful. Example: 4+ yellow/brown fine precipitate, 2+ small bipyramid crystals, clear drop, 3+ needle shaped crystals in 1+ white precipitate. One may also employ a standard numerical scoring scheme (Clear = 0, Precipitate = 1, Crystal = 10, etc). Figure 5 shows typical examples of what one might observe in a crystallization experiment.

### Interpretation of the Results

1. Fewer nucleation sites.
2. An increase in the amount of time required to see crystals using microscopy (i.e. slower growth rate).
3. In some cases, different habit, or morphology.
4. In some cases, reduced sedimentation.
5. In some cases, larger, fewer crystals.

### Optimization

1. Change your protein concentration. If there are too many nucleation sites, reduce protein concentration. No nucleation, increase protein concentration. Be patient as the equilibration will require up to 4 to 8 times longer than identical conditions with out the gel. One may alter these growth times by increasing the sample concentration/precipitant concentration to increase the rate of growth and decreasing the value of these parameters to slow growth.

2. Alter your sample / gel / precipitant ratio.

3. Consider the influence of other variables such as temperature, precipitant type & concentration, the use of ligands or metals, detergents, and the many other variables which could influence crystal growth.

### References and Readings

1. Garcia-Ruiz, J.M., Moreno, A., Viedma, C., & Coll, M. (1993) *Mat. Res. Bull.*, Vol. 28, pp. 541-546.
2. Garcia-Ruiz J.M. . (1991) *Key Engineering Material Vols. 58*, pp. 87-106.
3. McPherson, A. (1985) *Methods in Enzymology*, 14,112.

# Silica Hydrogel Kit™

HAMPTON  
RESEARCH

Solutions for Crystal Growth

## User Guide

3

4. Provost, K. & Robert, M.C. (1991) *J. Crystal Growth*, 110, 258.
5. Robert, M.C. & Lefaucheu, F. (1988) *J. Crystal Growth*, 90, 358.
6. Robert, M.C., Provost, K., & Lefaucheu, F., *Crystallization of Nucleic Acids and Proteins, A Practical Approach*, Oxford University Press, 127-143, 1992.

### Related Products

- HR3-158** Cryschem Plate - 24 plate case  
**HR3-160** Cryschem Plate - 100 plate case
- HR3-211** 22 mm Plain Square cover slide- 1 oz case (~100 slides)  
**HR3-213** 22 mm Plain Square cover slide- 10 oz case (~1,000 slides)  
**HR3-215** 22 mm Siliconized Square cover slide - 1 oz case (~100 slides)  
**HR3-217** 22 mm Siliconized Square cover slide - 10 oz case (~100 slides)
- HR3-227** 22 mm Plain Circle cover slides - 1 oz case (~120 slides)  
**HR3-229** 22 mm Plain Circle cover slides - 10 oz case (~1,200 slides)  
**HR3-231** 22 mm Siliconized Circle cover slides - 1 oz pack (~100 slides)  
**HR3-233** 22 mm Siliconized Circle cover slides - 10 oz pack (~1,000 slides)
- HR3-310** Micro-Bridges - 100 pack sampler  
**HR3-312** Micro-Bridges - 400 pack  
**HR3-340** Micro-Bridges Polypropylene - 100 pack sampler  
**HR3-342** Micro-Bridges Polypropylene - 400 pack
- HR4-110** Paper Wicks 25 mm - Assorted (XXX Fine - X Fine) - 200 wicks  
**HR4-122** Paper Wicks 25 mm - XXX Fine - 200 wicks  
**HR4-112** Paper Wicks 25 mm - XX Fine - 200 wicks  
**HR4-114** Paper Wicks 25 mm - X Fine - 200 wicks  
**HR4-116** Paper Wicks 25 mm - Fine - 200 wicks  
**HR4-118** Paper Wicks 25 mm - Medium - 200 wicks  
**HR4-120** Paper Wicks 25 mm - Coarse - 200 wicks
- HR4-211** Paper Wicks 55 mm - X Fine Long - 100 wicks  
**HR4-213** Paper Wicks 55 mm - Medium Long - 100 wicks  
**HR4-215** Paper Wicks 55 mm - Coarse Long - 100 wicks
- HR4-310** Red Sticky Wax - 80 wax ropes  
**HR4-312** Beeswax Sticks - 7 sticks  
**HR4-328** Capillary Wax - 40 gram pack (approximate)  
**HR4-326** Four Color Mounting Clay - 16 oz pack (4 oz each, blue / yellow / red / green)  
**HR4-342** Wax Pen - 1 each  
**HR4-344** Wax Pen Replacement Tip- 3 pack
- HR3-511** 1.88" Wide CrystalClear Sealing Tape & Dispenser  
**HR4-506** 3" Wide CrystalClear Sealing Tape  
**HR4-508** 4" Wide CrystalClear Sealing Tape  
**HR4-511** 2" Wide CrystalClear Sealing Tape  
**HR4-674** 1.88" Wide CrystalClear Sealing Tape, 3 pack  
**HR4-676** Dispenser for 1.88 inch wide tape, each, with one roll of tape
- HR4-334** Capillary Cutting Stone - 1 each
- HR6-104** 0.1 mm Glass #50 Capillaries - 25 pack  
**HR6-106** 0.2 mm Glass #50 Capillaries - 25 pack  
**HR6-108** 0.3 mm Glass #50 Capillaries - 25 pack  
**HR6-110** 0.4 mm Glass #50 Capillaries - 25 pack  
**HR6-112** 0.5 mm Glass #50 Capillaries - 25 pack  
**HR6-114** 0.6 mm Glass #50 Capillaries - 25 pack  
**HR6-116** 0.7 mm Glass #50 Capillaries - 25 pack  
**HR6-118** 0.8 mm Glass #50 Capillaries - 25 pack  
**HR6-120** 0.9 mm Glass #50 Capillaries - 25 pack  
**HR6-122** 1.0 mm Glass #50 Capillaries - 25 pack  
**HR6-124** 1.5 mm Glass #50 Capillaries - 25 pack  
**HR6-126** 2.0 mm Glass #50 Capillaries - 25 pack
- HR6-128** 0.1 mm Quartz Capillaries - 25 pack  
**HR6-130** 0.2 mm Quartz Capillaries - 25 pack  
**HR6-132** 0.3 mm Quartz Capillaries - 25 pack  
**HR6-134** 0.4 mm Quartz Capillaries - 25 pack  
**HR6-136** 0.5 mm Quartz Capillaries - 25 pack  
**HR6-138** 0.6 mm Quartz Capillaries - 25 pack  
**HR6-140** 0.7 mm Quartz Capillaries - 25 pack  
**HR6-142** 0.8 mm Quartz Capillaries - 25 pack  
**HR6-144** 0.9 mm Quartz Capillaries - 25 pack  
**HR6-146** 1.0 mm Quartz Capillaries - 25 pack  
**HR6-148** 1.5 mm Quartz Capillaries - 25 pack  
**HR6-150** 2.0 mm Quartz Capillaries - 25 pack  
**HR6-151** 2.5 mm Quartz Capillaries - 25 pack
- HR6-152** 0.1 mm Special Glass 10 Capillaries - 25 pack  
**HR6-154** 0.2 mm Special Glass 10 Capillaries - 25 pack  
**HR6-156** 0.3 mm Special Glass 10 Capillaries - 25 pack  
**HR6-158** 0.4 mm Special Glass 10 Capillaries - 25 pack  
**HR6-160** 0.5 mm Special Glass 10 Capillaries - 25 pack  
**HR6-162** 0.6 mm Special Glass 10 Capillaries - 25 pack  
**HR6-164** 0.7 mm Special Glass 10 Capillaries - 25 pack  
**HR6-166** 0.8 mm Special Glass 10 Capillaries - 25 pack  
**HR6-168** 0.9 mm Special Glass 10 Capillaries - 25 pack  
**HR6-170** 1.0 mm Special Glass 10 Capillaries - 25 pack  
**HR6-172** 1.5 mm Special Glass 10 Capillaries - 25 pack

# Silica Hydrogel Kit™



*Solutions for Crystal Growth*

<b>HR6-174</b>	2.0 mm Special Glass 10 Capillaries - 25 pack
<b>HR6-176</b>	0.1 mm Quartz Capillaries - 25 pack
<b>HR6-178</b>	0.2 mm Quartz Capillaries - 25 pack
<b>HR6-180</b>	0.3 mm Quartz Capillaries - 25 pack
<b>HR6-182</b>	0.4 mm Quartz Capillaries - 25 pack
<b>HR6-184</b>	0.5 mm Quartz Capillaries - 25 pack
<b>HR6-186</b>	0.6 mm Quartz Capillaries - 25 pack
<b>HR6-188</b>	0.7 mm Quartz Capillaries - 25 pack
<b>HR6-190</b>	0.8 mm Quartz Capillaries - 25 pack
<b>HR6-192</b>	0.9 mm Quartz Capillaries - 25 pack
<b>HR6-194</b>	1.0 mm Quartz Capillaries - 25 pack
<b>HR6-196</b>	1.5 mm Quartz Capillaries - 25 pack
<b>HR6-198</b>	2.0 mm Quartz Capillaries - 25 pack

## Technical Support

Inquiries regarding the Silica Hydrogel and general inquiries regarding crystallization are welcome. Please e-mail, fax, or telephone your request to Hampton Research. Fax and e-mail Technical Support are available 24 hours a day. Telephone technical support is available 8:00 a.m. to 5:00 p.m. USA Pacific Standard Time.

Hampton Research  
34 Journey  
Aliso Viejo, CA 92656-3317 U.S.A.  
Tel: (949) 425-1321 • Fax: (949) 425-1611  
Technical Support e-mail: [tech@hrmail.com](mailto:tech@hrmail.com)  
Website: [www.hamptonresearch.com](http://www.hamptonresearch.com)