

# PRODUCT DATA SHEET

**Product:** Anti-Fas Ligand, clone A11

Cat. No.: MC-105 (100 μg)

#### Specificity:

Fas Ligand (FasL) is a type II membrane protein and a member of the tumor necrosis factor family. When interacting with the Fas receptor, the trimeric FasL induces apoptosis. It is responsible for the activation-induced apoptosis of peripheral T cells and functions as an effector molecule for cytotoxic T cells. FasL is expressed in activated lymphocytes, Sertoli cells, and some tumor cells.

### Species Reactivity:

Mouse. Others not tested.

#### Ig Isotype:

Rat IgM

#### Immunogen:

Peptide corresponding to aa 196-220 of the mouse Fas Ligand protein

### Format:


100  $\mu$ L of 1 mg/mL antibody in PBS with 0.02% sodium azide. The antibody was purified from concentrated hybridoma tissue culture supernatant. Purity > 95% by SDS-PAGE,

### Storage:

Store at 4°C.

## Applications and Suggested Dilutions:

- Flow cytometry
- Immunohistochemistry: Excellent for this application.



Frozen Sections- Use 20  $\mu$ g/mL, incubate 1 hr at room temp. Method: Cryostat sections (6-8  $\mu$ m) were air dried, fixed in 4% paraformaldehyde in PBS and used immediately. Sections were incubated for 1 hr at RT with 20  $\mu$ g/mL of A11 antibody in PBS, 0.2% BSA, 0.05% Tween 20. After rinsing in PBS, a FITC-conjugated anti-rat IgM was added for 30 min, slides washed again in PBS and visualized using a fluorescence microscope.]

Paraffin-Embedded Sections- Dewax in xylene (2 x 10 min), EtOH 100% (2 x 10 min), block endogenous peroxidase (0.6%  $H_2O_2$  in Methanol 10 min), rehydrate through graded EtOH (100%, 95%, 70%), rinse in  $H_2O$ , unmask epitopes by heating slides in citrate buffer (0.01 M, pH 6.0) 2 x 5 min in microwave oven (650-750W), cool down to RT 20 min, wash 1x in PBS, incubate with A11 20  $\mu$ g/mL antibody, wash in PBS, incubate in anti-rat IgM HRP, wash in PBS, develop in appropriate substrate solution (DAB or AEC).

The optimal dilution for a specific application should be determined by the researcher.

#### Limitations:

For *in vitro* research use only. Not for use in diagnostics or in humans.

## Warranty:

No warranties, expressed or implied, are made regarding the use of this product. KAMIYA BIOMEDICAL COMPANY is not liable for any damage, personal injury, or economic loss caused by this product.

