

Ludger

reconstituted vial briefly before use. Ensure that any glass, plasticware or solvents used are free of glycosidases and environmental carbohydrates.

Minimise exposure to elevated temperatures or extremes of pH. High temperatures and low pH will cause desialylation. High pH will cause epimerisation of the reducing terminus GlcNAc.

Safety: This product is non-hazardous and has been purified from natural sources certified to be free of all hazardous material including pathogenic biological agents.

For research use only. Not for human or drug use

Related Products

Ludger Cat. No.	Description
CN-A2-x	A2 Glycan (di-sialylated parent of A1 glycan)
CN-NA2-x	NA2 Glycan (degalactosylated derivative of A1 glycan)
CN-NGA2-x	NGA2 Glycan (a substructure of NA2 glycan)
CN-M3N2-x	M3N2 Glycan (a substructure of NGA2 glycan)

Warranties and liabilities

Ludger warrants that the above product conforms to the attached analytical documents. Should the product fail for reasons other than through misuse Ludger will, at its option, replace free of charge or refund the purchase price. This warranty is exclusive and Ludger makes no other warrants, expressed or implied, including any implied conditions or warranties of merchantability or fitness for any particular purpose.

Ludger shall not be liable for any incidental, consequential or contingent damages.

This product is intended for *in vitro* research only.

Document # 'CN-A1-Guide', revision 1.2



Certificate of Analysis

A1 Glycan

Cat. #: CN-A1-10U

Lot #: A5AP-01

Size : 10 µg

Purity: > 90% pure as assessed by a combination of NMR and HPLC (see Fig 1).




Figure 1 : HPAE-PAD HPLC Profile of A1 Glycan (Cat. No.CN-A1-10U, Lot No. A5AP-01)