

Ludger™

NGA3 Glycan

Cat. No. CN-NGA3-x (where x denotes pack size)

Structure

$$\begin{array}{c} \text{GlcNAc}\beta 1 - 2 \, \text{Man} \, \alpha 1 \\ \\ \text{GlcNAc}\beta 1 - 4 \, \text{GlcNAc} \, \beta 1 - 4 \, \text{GlcNAc} \, \beta 1 - 4 \, \text{GlcNAc} \, \beta 1 \\ \\ \text{GlcNAc}\beta 1 - 2 \, \text{Man} \, \alpha 1 \end{array}$$

Synonyms: NGA3 N-linked oligosaccharide.

Description: Asialo-, agalacto-, tri-antennary complex-type N-glycan (oligosaccharide). NGA3 is a

substructure of NA3 glycan.

Sources : NGA3 glyan is a substructure of triantennary glycans found on several mammalian

glycoproteins including bovine serum fetuin (bSF). This product is typically purified from the oligosaccharide pool released from bSF by hydrazinolysis using a combination

of HPLC and glycosidase digestion.

Form: Dry. Dried by centrifugal evaporation from an aqueous solution.

Molecular Weight: 1521

Purity: > 90% pure as assessed by a combination of ¹H-NMR and HPLC.

Storage: Refridgerate (-20°C) both before and after dissolution. This product is stable for at least

5 years as supplied.

Shipping: The product can be shipped at ambient when dry. After dissolution, ship on dry ice.

Handling: Allow the unopened vial to reach ambient temperature and tap unopened on a solid

surface to ensure that most of the lyophilized material is at the bottom of the vial.

Gently remove the cap, add the desired volume of reconstitution medium, re-cap and mix thoroughly to bring all the oligosaccharide into solution. For maximal recovery of





oligosaccharide, ensure that the cap lining is also rinsed and centrifuge the reconstituted vial briefly before use. Ensure that any glass, plasticware or solvents

used are free of glycosidases and environmental carbohydrates.

Safety: This product is non-hazardous and has been purified from natural sources certified to

be free of all hazardous material including pathogenic biological agents.

For research use only. Not for human or drug use

Related Products

Ludger Description

Cat. No.

CN-A3-x A3 Glycan (trisialylated parent of NA3 glycan)

CN-NA3-x NA3 Glycan (the parent structure of NGA3 glycan)

CN-M3N2-x M3N2 Glycan (a substructure of NGA3 glycan)


Warranties and liabilities

Ludger warrants that the above product conforms to the attached analytical documents. Should the product fail for reasons other than through misuse Ludger will, at its option, replace free of charge or refund the purchase price. This warranty is exclusive and Ludger makes no other warrants, expressed or implied, including any implied conditions or warranties of merchantability or fitness for any particular purpose.

Ludger shall not be liable for any incidental, consequential or contingent damages.

This product is intended for in vitro research only.

Revised: 07.June.2001



Ludger™

Certificate of Analysis

NGA3 Glycan

Cat. #: CN-NGA3-20U

Lot #: A15H-02

Size : 20 μg

Purity: > 90% pure as assessed by a combination of ¹H-NMR (see Fig 1) and HPLC.




Figure 1: 500 MHz 1H-NMR of NGA3 Glycan (Cat. No.CN-NGA3-20U, Lot No. A15H-02)