

Product Information Sheet

Recombinant Murine PDGF-BB

Catalog# 315-18

Source: *E.coli*

Description: PDGFs are disulfide-linked dimers consisting of two 12.0-13.5 kDa polypeptide chains, designated PDGF-A and PDGF-B chains. The three naturally occurring PDGFs; PDGF-AA, PDGF-BB and PDGF-AB, are potent mitogens for a variety of cell types including smooth muscle cells, connective tissue cells, bone and cartilage cells, and some blood cells. The PDGFs are stored in platelet α -granules and are released upon platelet activation. The PDGFs are involved in a number of biological processes, including hyperplasia, chemotaxis, embryonic neuron development, and respiratory tubule epithelial cell development. Two distinct signaling receptors used by PDGFs have been identified and named PDGFR- α and PDGFR- β . PDGFR- α is high-affinity receptor for each of the three PDGF forms. On the other hand, PDGFR- β interacts with only PDGF-BB and PDGF-AB. Recombinant murine PDGF-BB is a 24.4 kDa disulfide-linked homodimer of two B chains (218 total amino acids).

Synonyms: Platelet-Derived Growth Factor-BB, Glioma-derived growth factor (GDGF), Osteosarcoma-derived Growth Factor (ODGF)

Sequence: SLGSLAAAEP AVIAECKTRT EVFQISRNL I DRTNANFLVW PPCVEVQRCS GCCNNRNVQC RASQVQMRPV QVRKIEIVRK
KPIFKKATVT LEDHLACKCE TIVTPRPVT

Authenticity: Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Purity: $\geq 98\%$ by SDS-PAGE gel and HPLC analyses.

Endotoxin: Endotoxin level is < 0.1 ng/ μ g of protein (< 1 EU/ μ g).

Protein Content: Verified by UV Spectroscopy and/or SDS-PAGE gel.

Biological activity: Determined by the dose-dependent stimulation of the proliferation of Balb/c 3T3 cells. The expected ED₅₀ for this effect is 1.0-2.0 ng/ml.

For a list of references please visit our website at www.peprotech.com

Country of Origin: USA

Lot-specific information is not available on Product Information Sheet.
See Certificate of Analysis for details.

Usage: For Research Use Only. Not for use in diagnostic or therapeutic procedures.

Updated 07/15/2013

Storage/Stability:

Product Form	Temperature	Storage Time
Lyophilized	-20°C to -80°C	See expiration date
Lyophilized	4°C	6 months
Lyophilized	RT	1 month
Diluted as per CoA	-20°C to -80°C	12 months
Diluted as per CoA	2°C to 8°C	1 week
**Avoid repeated freeze-thaw cycles.		