



Mouse Anti-Human CD8

Cat. No.	Form	Quantity
9536-01	Purified (UNLB) Antibody	0.1 mg
9536-02	Fluorescein (FITC) Conjugate	100 tests
9536-08	Biotin (BIOT) Conjugate	100 tests
9536-09	R-phycoerythrin (R-PE) Conjugate	100 tests
9536-10	R-phycoerythrin-Texas Red [®] (R-PE-TXRD) Conjugate	100 tests
9536-11	Allophycocyanin (APC) Conjugate	100 tests
9536-13	*Spectral Red [™] (SPRD) Conjugate	100 tests
9536-14	Low Endotoxin, Azide-Free (LE/AF)	0.1 mg
9536-15	**Cyanine 5 (CY [™] 5) Conjugate	100 tests
9536-16	**R-phycoerythrin-Cyanine 5.5 (R-PE-CY [™] 5.5) Conjugate	100 tests
9536-17	**R-phycoerythrin-Cyanine 7 (R-PE-CY [™] 7) Conjugate	100 tests
9536-18	** Allophycocyanin -Cyanine 5.5 (APC-CY [™] 5.5) Conjugate	100 tests
9536-19	** Allophycocyanin -Cyanine 7 (APC-CY [™] 7) Conjugate	100 tests

DESCRIPTION

Clone RFT-8
Ig Isotype Mouse IgG₁
Specificity Human CD8

CD8 is a 32 kDa transmembrane glycoprotein expressed as either a CD8 $\alpha\alpha$ homodimer or CD8 $\alpha\beta$ heterodimer. It is expressed on the “cytotoxic/suppressor” subpopulation of peripheral T cells. CD8 functions primarily as a coreceptor with MHC Class I-restricted TCR’s in antigen recognition.¹⁻²

RESEARCH APPLICATIONS

- Flow cytometry
- Immunohistochemistry (frozen sections)

CHARACTERIZATION

To insure acceptable performance, each batch of product is tested by flow cytometry to conform to the characteristics of a standard reference reagent.

WORKING DILUTIONS

Flow Cytometry:	Purified antibody	≤ 1 μg/10 ⁶ cells
	Fluorescein conjugate	10 μL/10 ⁶ cells
	Biotin conjugate	10 μL/10 ⁶ cells
	R-phycoerythrin conjugate	10 μL/10 ⁶ cells
	R-phycoerythrin conjugate-Texas Red [®]	10 μL/10 ⁶ cells
	Allophycocyanin conjugate	10 μL/10 ⁶ cells
	Spectral Red [™] conjugate	10 μL/10 ⁶ cells
	Cyanine 5 conjugate	10 μL/10 ⁶ cells
	R-phycoerythrin-Cyanine 5.5 conjugate	10 μL/10 ⁶ cells
	R-phycoerythrin-Cyanine 7 conjugate	10 μL/10 ⁶ cells
	Allophycocyanin-Cyanine 5.5 conjugate	10 μL/10 ⁶ cells
	Allophycocyanin-Cyanine 7 conjugate	10 μL/10 ⁶ cells

Other Applications: Since applications vary, investigators are advised to determine the optimum working dilution of the product that is appropriate for a specific use.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

HANDLING AND STORAGE

- The purified (UNLB) antibody is supplied as 0.1 mg of purified immunoglobulin in 1.0 mL of 100 mM borate buffered saline, pH 8.0. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) and Cyanine 5 (CYTM5) conjugates are supplied as 100 tests in 1.0 mL PBS/NaN₃. Store at 2-8°C.
- The biotin (BIOT) conjugate is supplied as 100 tests in 1.0 mL PBS/NaN₃. Store at 2-8°C.
- The R-phycoerythrin (R-PE) and allophycocyanin (APC) conjugates are supplied as 100 tests in 1.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The Spectral RedTM (SPRD), R-phycoerythrin -Texas Red[®] (R-PE-TXRD), R-phycoerythrin-Cyanine 7 (R-PE-CYTM7), R-phycoerythrin-Cyanine 5.5 (R-PE-CYTM5.5), Allophycocyanin-Cyanine 5.5 (APC-CYTM5.5), and Allophycocyanin-Cyanine 7 (APC-CYTM7), conjugates are supplied as 100 tests in 1.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The low endotoxin, azide-free (LE/AF) antibody is supplied as 0.5 mg purified immunoglobulin in 0.2 mL of PBS. **Aliquot and store at or below -20°C.**
- Protect conjugated forms from light. Aliquot and freeze the low endotoxin, azide-free product at -20°C immediately upon receipt. Each reagent is stable for the period shown on the bottle label if stored as directed.

WARNING

Reagents contain sodium azide which is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

REFERENCES

1. Barclay, A.N., M.H. Brown, S.K.A. Law, A.J. McKnight, M.G. Tomlinson, and P.A. van der Merwe, eds. 1997. *The Leukocyte Antigens Facts Book, 2nd Edition*, CD8 Section, Academic Press, New York, p. 149.
2. Zamoyska, R. 1994. *Immunity* 1:243.

Texas Red[®] is a registered trademark of Molecular Probes, Inc.

FACScanTM is a registered trademark of Becton Dickinson Immunocytometry Systems, Inc., San Jose, CA.

Spectral RedTM is a registered trademark of Southern Biotechnology Associates, Inc.

*Spectral RedTM is a PE/CyTM5 tandem conjugate. **CyTM5, CyTM7 and CyTM5.5 are trademarks of Amersham Biosciences Corp. CyTM5, CyTM7 and CyTM5.5 are for non-commercial research use only, not for therapeutic or in vivo applications. Other use needs license from Amersham Biosciences Corp., under U.S. Patent Nos. 4,981,977 and 5,268,486 and other patents pending. This material (or portions of this material) is subject to proprietary rights of Amersham Biosciences Corp. and Carnegie Mellon University and made and sold under license from Amersham Biosciences Corp.. This product is licensed for sale only for research. It is not licensed for any other use. There is no implied license hereunder for any commercial use. Commercial Use shall include; 1. sale, lease, license or other transfer of the material or any material derived or produced from it. 2. sale, lease, license or other grant of rights to use this material or any material derived or produced from it. 3. Use of this material to perform services for a fee for third parties. If you require a commercial license to use this material and do not have one, return this material, unopened to Southern Biotechnology, 160A Oxmoor Blvd, Birmingham, AL 35209, USA and any money paid for the material will be refunded.