



Instructions For Use

VGS-IFU

Rev. Date: Jan. 29, 2012

Revision: 3

Page 1 of 1

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Tel. (435) 755-9848 - Fax (435) 755-0015 - www.scytek.com

Van Gieson's Solution

Description: The Van Gieson's Solution is intended for use in the histological visualization of collagen in tissue

sections. This reagent is used in many special stain procedures including the Colloidal Iron Stain Kit and

the Elastic Stain Kit (Modified Verhoff's).

Collagen: Red Muscle Fibers: Yellow Cytoplasm: Yellow

Uses/Limitations: Not to be taken internally.

For In-Vitro Diagnostic use only.


Histological applications.

Do not use if reagents become cloudy. Do not use past expiration date. Use caution when handling reagents.

Non-Sterile.

Control Tissue: Lung

Uterus Muscle Kidney



Availability: <u>Item #</u> <u>Volume</u>

VGS125 125 ml VGS500 500 ml VGS999 1000 ml

Storage: Store at 18-25℃.

Precautions: Avoid contact with skin and eyes.

Harmful if swallowed.

Follow all Federal, State, and local regulations regarding disposal.

Procedure:

- 1. Deparaffinize sections if necessary and hydrate to distilled water.
- 2. Apply adequate Van Gieson's Solution to completely cover tissue section and incubate for 1 minute.
- Rinse slide using absolute alcohol.
- 4. Dehydrate in 2 changes of absolute alcohol, clear, and mount in synthetic resin.

U.S.A.

Storage: 18° C



ScyTek Laboratories, Inc. 205 South 600 West Logan, UT 84321

CE IND

EC REP EmergoEurope (31)(0) 70 345-8570 Molsnstraat 15 2513 BH Hague, The Netherlands

Doc: IFU-Template18-25rev2