

DNA STAT

DNA STAT-60™

SINGLE REAGENT FOR RAPID DNA ISOLATION

Cat No. TL-4200

For in vitro research use

TEL-TEST "B"
Bulletin No. 9

1. INTRODUCTION:

DNA STAT-60, a single monophasic reagent containing a chaotropic cell disrupter and a non-corrosive phenol free extraction reagent replaces cumbersome, time, and labor intensive methods of genomic DNA isolation. Following tissue or cell homogenization in the DNA STAT-60, and after the addition of chloroform, the homogenate separates into two phases: the aqueous phase and organic phase. The DNA remains in the aqueous phase while RNA and the other cellular components, including proteins are preferentially partitioned in the organic phase or interface. The DNA STAT-60 method does not require ultracentrifugation, and can be completed in under 1 hour. The DNA STAT-60 isolates high molecular weight genomic DNA from samples of human, animal, plant, yeast, and bacterial origin and is particularly well suited for the simultaneous processing of multiple samples.

RNA and proteins are sequestered in organic and interphase; DNA remains in upper aqueous phase.

Protocol can be completed in 60 minutes.

Single extraction step replaced cumbersome, time, and labor intensive steps requiring proteinase K, phenol, or solid supports.

Single monophasic reagent with extended shelf life.

Does not contain phenol or hazardous organics.

Isolates genomic DNA from samples of human, animal, plant, yeast and bacterial origin. Can be used to back extract DNA from "SINGLE STEP METHOD" RNA preps (i.e., RNAzol, RNAzol B, RNA STAT-60)

2. REAGENTS SUPPLIES:

DNA STAT-60™ 50ml, 100ml, or 200ml bottle containing clear solution of DNA STAT-60™.

PREPARATION: Ready to use.

STORAGE: Refrigerate at 2-8°C. Protect from exposure to light.

STABILITY: 6 months. Refer to expiration date stamped on label.

3. REAGENTS REQUIRED, BUT NOT SUPPLIED:

Chloroform (ACS grade)
Isopropanol (ACS grade)
Ethanol (ACS grade)

4. PROTOCOL:

DNA isolation by the DNA STAT-60™ method includes the following steps:

1. Homogenization DNA STAT-60™
 (1ml per 50-100mg tissue, or 5-10 x 10⁶ cells.
2. DNA Extraction 1 vol. of homogenate +
 0.2 vol. of chloroform
3. DNA Precipitation 0.5 vol. of isopropanol
4. DNA Wash 75% ethanol

Unless stated otherwise the procedure is carried out at room temperature.

Continued on other side

4.1 HOMOGENIZATION:

A. TISSUES:

Homogenize tissues in the DNA STAT-60 (1ml/65mg issue) in a glass-teflon or polytron homogenizer.

B. CELLS:

Cells grown in monolayer are lysed directly in a culture dish by adding DNA STAT-60 (1ml/3.5cm petri dish) and passing cell lysate several times through a pipette. Cells grown in suspension are sedimented then lysed in a DNA STAT-60 (1ml per 5×10^6 cell) by repetitive pipetting.

4.2 DNA EXTRACTION:

Following homogenization add 0.2ml of chloroform per 1ml of DNA STAT-60, cover the sample tightly, shake vigorously for 15 seconds and let it stay at room temperature for 2-3 minutes. Centrifuge the homogenate at 12,000g(max) for 15 minutes at 4°C. Following centrifugation the homogenate separates into two phases: a lower organic phase and the upper aqueous phase. DNA remains in the aqueous phase whereas RNA and proteins are in interphase and organic phase.

4.3 DNA PRECIPITATION

Transfer the aqueous phase to a fresh tube and mix with isopropanol. Add 0.5ml of isopropanol per 1ml of the DNA STAT-60 used for homogenization. Store samples at room temperature for 5-10 minutes and centrifuge at 12,000g(max) for 10 minutes at 4°C. The DNA precipitate forms a small clear to white pellet at the bottom of the tube.

4.4 DNA WASH

Remove supernatant and wash the DNA pellet once with 75% ethanol by vortexing and subsequent centrifugation at 7,500g(max) for 5 minutes at 4°C. Add at least 1ml of 75% ethanol per 1ml of the DNA STAT-60 used for the initial homogenization.

At the end of the procedure, dry the DNA pellet briefly by air drying or in a vacuum (5-10 min.). Dissolve the DNA pellet in water or in 1mM EDTA, pH 7. Vortex or pass the pellet a few times through a pipette tip. An incubation for 10-15 minutes at 55-60°C may be required to dissolve DNA samples.

5. RECOVERING DNA FROM "SINGLE STEP METHOD" (RNAzol, RNAzol B, RNA STAT-60 PREPS.

5.1 DNA REVERSE EXTRACTION

Remove aqueous layer containing RNA. Add 800ul of DNA STAT-60 reagent per 1 ml of RNAzol, RNAzol B or RNA STAT-60 used for the initial homogenization. Add 0.2ml of chloroform per 1ml of DNA STAT-60, shake vigorously for 15 seconds and let it stay at room temperature for 2-3 minutes. Centrifuge the homogenate at 1000g(min)-12,000g(max) for 15 minutes 4°C. Following centrifugation the homogenate separates into two phases: a lower organic phase and the upper aqueous phase. DNA remains in the aqueous phase whereas RNA and proteins are in the interphase and organic phase.

5.2 DNA PRECIPITATION

Follow procedure as outlined in section 3.3.

5.3 DNA WASH

Follow procedure as outlined in section 3.4.

6. SPECIAL HANDLING PRECAUTIONS

The DNA STAT-60 contains an irritant (Guanidinium Salts). Can be fatal. When working with DNA STAT-60 use gloves and eye protection (shield, safety goggles). Do not get on skin or clothing. Avoid breathing vapor. Read warning note on bottle.

REFERENCES

1. Chomczynski, P. and Sacchi, 1987. Single-Step Method of RNA Isolation by Acid Guanidinium Thiocyanate-Phenol-Chloroform Extraction. *Biotechniques* 8:148-149.
2. Maniatis, T., E.T. Fritsch and J. Sambrook. 1982. *Molecular Cloning: A Laboratory Manual*.
3. Winberg, G. 1991. A Rapid Method for Preparing DNA from Blood, Suited for PCR Screening of Transgenes in Mice. *PCR Methods Applic.* 1:72-74.

DNA STAT-60™

CAT. NO. TL-4200 50ml

CAT. NO. TL-4210 100ml

CAT. NO. TL-4220 200ml

DNA STAT-60™ is a trademark of Tel-Test "B", Inc.

TEL-TEST "B", INC.

P. O. Box 1421 · 1511 County Road 129 · Friendswood, Tx 77546
Customer Service (713) 482-2672 · Fax (713) 482-1070
Ordering (outside Texas) 1-800-631-0600