

Product Insert

AimPlex™ Analyte Kit

Mouse Chemokine 7-Plex (96 Tests)

PN: C283308

COMPONENTS

1x Ab-conjugated beads (premixed 8-plex). PN: C283308A. One bottle containing 5 mL of the following AimPlex antibody-bead conjugates:

S4P5 - anti-mouse IP-10
S4P9 - anti-mouse KC
S4P10 - anti-mouse Eotaxin
S5P3 - anti-mouse MCP-1
S5P7 - anti-mouse MIG
S5P8 - anti-mouse RANTES
S5P10 - anti-mouse MCP-3

2x Biotin-detection Ab (premixed 8-plex). PN: C283308B. One vial containing 1.5 mL of the following biotinylated antibodies.

Biotin - anti-mouse IP-10
Biotin - anti-mouse KC
Biotin - anti-mouse Eotaxin
Biotin - anti-mouse MCP-1
Biotin - anti-mouse MIG
Biotin - anti-mouse RANTES
Biotin - anti-mouse MCP-3

2x Mouse/Rat dAb Diluent. PN: 07MRDX21. One vial containing 1.5 mL of biotin detection antibody diluent

Lyophilized Standard Mix-Mouse Chemokine 7-Plex. PN: MG19307. One vial containing lyophilized recombinant mouse Eotaxin, IP-10, KC, MCP-1, MCP-3, MIG, RANTES.

STORAGE: 2-8°C in the dark.

IMPORTANT: Sodium azide forms explosive compounds with heavy metals. These products contain <0.05% (w/w) azide which with repeated contact with lead and copper commonly found in plumbing drains may result in the buildup of shock sensitive compounds. Dispose in accordance with regulations from your institute.

APPLICATION: Optimal antibody pairs and antigen standards for assaying mouse Inflammation 8-Plex Panel. To be used in conjunction with the AimPlex Basic Kit for Premixed Panels (PN: U100005) and a diluent kit. Refer to the AimPlex Multiplex Immunoassay User Manual and kit inserts for the assay procedure.

For Research Use Only. Not for use in diagnostic procedures.

Assay Specifications:

Sample types: Cell culture supernatant, serum, plasma, bodily fluid and tissue/cell lysate

	Sensitivity (LOD, pg/mL)	LLOQ (pg/mL)	ULOQ (pg/mL)
Mouse Eotaxin	< 3	< 10	> 2000
Mouse IP-10	< 3	< 10	> 5000
Mouse KC	< 1	< 2	> 2000
Mouse MCP-1	< 1	< 2	> 2000
Mouse MCP-3	< 2	< 5	> 5000
Mouse MIG	< 2	< 5	> 5000
Mouse RANTES	< 2	< 5	> 5000

Standard dose recovery: 70-130%

Intra-assay CV: < 10%

Inter-assay CV: < 20%

Cross-reactivity of analytes in the panel: Negligible

Sample volume: 15 µL/test

Please contact Customer Service at 626-817-6366 with any questions or comments.

Thank you for your business!