

Anti-4-Hydroxy-2-nonenal (4-HNE) *Monoclonal Antibody (HNEJ-2)*

Code: **MHN-100P**
(100 μ g of IgG, Lyophilized powder)
MHN-020P
(20 μ g of IgG, Lyophilized powder)

Source: **Mouse**

Immunogen: 4-Hydroxy-2-nonenal (4-HNE)-modified keyhole limpet hemocyanine.

Subclass: IgG1 (κ)

Applications: Immunohistochemistry [see ref.1]
(Recommended concentration: 25 μ g/mL IgG)

Western blotting [see ref.2]
(Recommended concentration: 15 μ g/mL IgG)

Reconstitution: MHN-100P: Dissolve in 1mL of distilled water.
MHN-020P: Dissolve in 1mL of distilled water.

Buffer Concentration: 100 μ g/mL IgG in 50mM Tris-buffered saline (TBS).

Specificity: This antibody show almost negligible reactivity with proteins that were treated with other aldehydes, such as 2-nonenal, 2-hexenal, 1-hexanal, 4-hydroxy-2-hexenal, formaldehyde, or glutaraldehyde.
By inhibition test, this antibody has a much higher affinity for the 4-HNE-histidine adduct than 4-HNE-lysine or 4-HNE cysteine adduct.

Storage: Store at less than -20°C.
Avoid repeated freeze & thaw after reconstitution.
For short term storage or transport, storage at 4°C is acceptable.

Stability: 5 years at -20°C.

References: 1) T.Tanaka *et al.* *Laboratory Investigation* 77(2), p145-155 (1997)
2) S.Toyokuni *et al.* *FEBS Letters* 359, p189-191 (1995)

For research use only, not for diagnostic use.

Manufacturer



JAPAN INSTITUTE FOR THE CONTROL OF AGING (JaICA)



COSMO BIO Co., LTD.
Inspiration for Life Science

TOYO EKIMAE BLDG, 2-20, TOYO 2CHOME, KOTO-KU, TOKYO 135-0016, JAPAN
TEL: +81-3-5632-9617 FAX: +81-3-5632-9618 e-mail: export@cosmobio.co.jp

www.cosmobio.co.jp